

AN EFFECTIVE CRITERION FOR EULERIAN MULTIZETA VALUES IN POSITIVE CHARACTERISTIC

CHIEH-YU CHANG, MATTHEW A. PAPANIKOLAS, AND JING YU

ABSTRACT. Characteristic p multizeta values were initially studied by Thakur, who defined them as analogues of classical multiple zeta values of Euler. In the present paper we establish an effective criterion for Eulerian multizeta values, which characterizes when a multizeta value is a rational multiple of a power of the Carlitz period. The resulting “ t -motivic” algorithm can tell whether any given multizeta value is Eulerian or not. We also prove that if $\zeta_A(s_1, \dots, s_r)$ is Eulerian, then $\zeta_A(s_2, \dots, s_r)$ has to be Eulerian. When $r = 2$, this was conjectured (and later on conjectured for arbitrary r) by Lara Rodríguez and Thakur for the zeta-like case from numerical data. Our methods apply equally well to values of Carlitz multiple polylogarithms at algebraic points and zeta-like multizeta values.

1. INTRODUCTION

In this paper we provide an effective criterion to determine when multizeta values in positive characteristic are Eulerian. Our study is motivated by the celebrated formula of Euler on special values of the Riemann zeta function at even positive integers: for $m \in \mathbb{N}$,

$$\zeta(2m) = \frac{-B_{2m} (2\pi\sqrt{-1})^{2m}}{2(2m)!},$$

where $B_{2m} \in \mathbb{Q}$ are Bernoulli numbers. In particular, we have $\zeta(2m)/(2\pi\sqrt{-1})^{2m} \in \mathbb{Q}$ for $m \in \mathbb{N}$. For an integer $n > 1$, Euler’s formula implies (trivially, since $\zeta(n)$ is real) that $\zeta(n)/(2\pi\sqrt{-1})^n$ is rational if and only if n is even.

Multiple zeta values (henceforth abbreviated MZV’s), initially studied by Euler as generalizations of special zeta values, are defined by the reciprocal power sums

$$\zeta(s_1, \dots, s_r) = \sum_{n_1 > \dots > n_r \geq 1} \frac{1}{n_1^{s_1} \cdots n_r^{s_r}},$$

where s_1, \dots, s_r are positive integers with $s_1 \geq 2$. Here r is called the depth and $w := \sum_{i=1}^r s_i$ is called the weight of $\zeta(s_1, \dots, s_r)$. We call $\zeta(s_1, \dots, s_r)$ *Eulerian* if the ratio $\zeta(s_1, \dots, s_r)/(2\pi\sqrt{-1})^w$ is rational (see [To4]). It is a natural question to ask if there is a criterion for determining which MZV’s of depths at least 2 are Eulerian.

Date: November 1, 2014.

2010 Mathematics Subject Classification. Primary 11R58, 11J93; Secondary 11G09, 11M32, 11M38.

Key words and phrases. Multizeta values, Eulerian, Carlitz tensor powers, Carlitz polylogarithms, Anderson-Thakur polynomials.

The first author was partially supported by a Golden-Jade fellowship of the Kenda Foundation, NCTS, and MOST Grant 102-2115-M-007-013-MY5. The second author was partially supported by NSF Grant DMS-1200577. The third author was partially supported by MOST Grant 102-2119-M-002-002.

Let A be the polynomial ring in the variable θ over a finite field \mathbb{F}_q with quotient field k . Let A_+ be the set of monic polynomials in A and consider the series, for $n \in \mathbb{N}$,

$$\zeta_A(n) := \sum_{a \in A_+} \frac{1}{a^n} \in \mathbb{F}_q\left(\left(\frac{1}{\theta}\right)\right).$$

These values, called Carlitz zeta values, are analogues of classical special zeta values. We note that in this non-archimedean situation the series $\zeta_A(1)$ does converge in $\mathbb{F}_q\left(\left(\frac{1}{\theta}\right)\right)$. Let \mathbf{C} be the Carlitz module and $\tilde{\pi}$ be a fundamental period of \mathbf{C} . Recall that in the function field setting \mathbf{C} plays the role of the multiplicative group \mathbb{G}_m and $\tilde{\pi}$ plays the role of $2\pi\sqrt{-1}$. We denote by $\exp_{\mathbf{C}}(z) = \sum_{i \geq 0} z^{q^i} / D_i$ the Carlitz exponential function, and by $\Gamma_{n+1} \in A$ (for non-negative integers n) the Carlitz factorials (see §4.1 for definitions).

In [Ca35], Carlitz derived an analogue of Euler's formula. More precisely, we write

$$\frac{z}{\exp_{\mathbf{C}}(z)} = \sum_{n \geq 0} \frac{\text{BC}(n)}{\Gamma_{n+1}} z^n,$$

where $\text{BC}(n) \in k$ are called Bernoulli-Carlitz numbers (see [Go96]). Carlitz established the formula

$$(1.0.1) \quad \zeta_A(n) = \frac{\text{BC}(n)}{\Gamma_{n+1}} \tilde{\pi}^n$$

if $n \in \mathbb{N}$ is *even* (i.e., $(q-1)|n$). We note that $\tilde{\pi}^n \in \mathbb{F}_q\left(\left(\frac{1}{\theta}\right)\right)$ if and only if n is *even*, and so Carlitz's result implies that $\zeta_A(n)/\tilde{\pi}^n \in k$ if and only if n is *even*.

In [AT90], Anderson and Thakur related $\zeta_A(n)$ to the last coordinate of the logarithm of $\mathbf{C}^{\otimes n}$ (the n -th tensor power of the Carlitz module viewed as a Carlitz-Tate t -motive) at an explicitly constructed integral point Z_n (see §5.1.2). As a consequence, one has that the rationality of $\zeta_A(n)/\tilde{\pi}^n$ is equivalent to Z_n being $\mathbb{F}_q[t]$ -torsion. In this case, it is clearly described when \mathbf{v}_n is $\mathbb{F}_q[t]$ -torsion, and more precisely we have that \mathbf{v}_n is an $\mathbb{F}_q[t]$ -torsion point if and only if n is *even* (see [AT90, Prop. 1.11.2, Cor. 3.8.4] and [Yu91, Thm. 3.1]). On the other hand, Yu showed that the transcendence of $\zeta_A(n)/\tilde{\pi}^n$ is equivalent to \mathbf{v}_n being non-torsion (see [Yu91, Cor. 2.6]), whence deriving that $\zeta_A(n)/\tilde{\pi}^n$ is algebraic over k if and only if $\zeta_A(n)/\tilde{\pi}^n$ is in k .

For $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$, characteristic p multizeta values $\zeta_A(s_1, \dots, s_r)$, defined by Thakur [To4], are generalizations of Carlitz zeta values. We set

$$\zeta_A(s_1, \dots, s_r) := \sum \frac{1}{a_1^{s_1} \cdots a_r^{s_r}} \in \mathbb{F}_q\left(\left(\frac{1}{\theta}\right)\right),$$

where the sum is taken over r -tuples of monic polynomials a_1, \dots, a_r with $\deg a_1 > \cdots > \deg a_r$, r is called the depth and $w := s_1 + \cdots + s_r$ is the weight. These values are known to be non-vanishing by Thakur [To9a, Thm. 4]. As in the classical case, Thakur called $\zeta_A(s_1, \dots, s_r)$ *Eulerian* if the ratio $\zeta_A(s_1, \dots, s_r)/\tilde{\pi}^w$ is in k . We mention that one encounters here the Eulerian multizeta values such as $\zeta_A(q-1, (q-1)^2)$, or $\zeta_A(q-1, (q-1)q, \dots, (q-1)q^{r-1})$ (see [To9b, LRT14, Ch14]), as compared with the classical Eulerian values $\zeta(2m, 2m)$, $\zeta(2, 2, \dots, 2)$. In contrast to the classical story, we already know that these ratios $\zeta_A(\mathfrak{s})/\tilde{\pi}^w$ are either rational or transcendental over k . Indeed, by [C14, Cor. 2.3.3] we have that either $\zeta_A(\mathfrak{s})/\tilde{\pi}^w$ is in k or $\zeta_A(\mathfrak{s})$ and $\tilde{\pi}$ are algebraically independent over k , generalizing the depth one results of [Yu97, CY07]. However the

“irrationality” remains a subtle question, i.e. verifying that a given specific *even* weight multizeta value of depth $r > 1$ is not Eulerian.

The main result of the present paper (Theorem 6.1.1) is to give an effective criterion for Eulerian multizeta values of arbitrary depth. Inspired by Anderson-Thakur [AT09], for any r -tuple $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$ we first explicitly construct an abelian t -module E' defined over A , which is a higher dimensional analogue of a Drinfeld module introduced by Anderson [A86], and an integral point $\mathbf{v}_{\mathfrak{s}} \in E'(A)$ such that $\zeta_A(s_1, \dots, s_r)$ is Eulerian if and only if $\mathbf{v}_{\mathfrak{s}}$ is an $\mathbb{F}_q[t]$ -torsion point in $E'(A)$. Furthermore, whenever $\zeta_A(s_1, \dots, s_r)$ is Eulerian we find an explicit polynomial $a_{\mathfrak{s}} \in \mathbb{F}_q[t]$ that annihilates the integral point $\mathbf{v}_{\mathfrak{s}}$. This allows us to construct an algorithm for determining when a given MZV is Eulerian or non-Eulerian. Interesting observations come up from implementation of this algorithm which will be discussed in §6.

In the classical case, Brown [B12b, Thm. 3.3] gave a sufficient condition for Eulerian MZV's in terms of motivic multiple zeta values, which are functions defined on the motivic period torsor for the motivic Galois group of the mixed Tate motives over \mathbb{Z} , and whose images under the period map are the multiple zeta values in question. Given any $\zeta(s_1, \dots, s_r)$ with even weight N , if the corresponding motivic multiple zeta value $\zeta^m(s_1, \dots, s_r)$ is trivial under the operator $D_{<N}$ given in [B12b, (3.2)], then Brown proves that $\zeta(s_1, \dots, s_r)$ is Eulerian.

Overall it seems difficult to verify when our integral point $\mathbf{v}_{\mathfrak{s}}$ is an $\mathbb{F}_q[t]$ -torsion point in $E'(A)$ directly in terms of s_1, \dots, s_r alone. Even in the classical case to date there is no conjecture that describes Eulerian MZV's precisely in terms of s_1, \dots, s_r . However, as a notable consequence of the main result, we find that if $\zeta_A(s_1, \dots, s_r)$ is Eulerian, then the $r - 1$ MZV's, $\zeta_A(s_2, \dots, s_r), \dots, \zeta_A(s_r)$ are simultaneously Eulerian (see Corollary 4.2.3). For the classical MZV's, Brown's theorem on a sufficient condition for Eulerian MZV's implies that $\zeta(3, 1, \dots, 3, 1)$ is Eulerian (see [B12a, Rem. 4.8]). However, $\zeta(1)$ does not converge and so a naive analogue of the truncation result for the classical Eulerian MZV's does not make sense. It would be interesting to ask whether some analogue of the characteristic p truncation is nevertheless valid for the classical Eulerian MZV's without 1 occurring in the coordinates.

The methods of constructing t -modules together with specific integral points which are developed in this paper also enable us to investigate similar phenomena for zeta-like multizeta values. As defined by Thakur, $\zeta_A(s_1, \dots, s_r)$ is called *zeta-like* if the ratio

$$\zeta_A(s_1, \dots, s_r) / \zeta_A\left(\sum_{i=1}^r s_i\right)$$

is in k (equivalently it is algebraic over k by [C14, Thm. 2.3.2]). A criterion for zeta-like MZV's (see Theorem 5.3.6) is given in terms of $\mathbb{F}_q[t]$ -linear relations for the corresponding two integral points on our t -modules. Working out this criterion, we also deduce the fact that having $\zeta_A(s_1, \dots, s_r)$ zeta-like implies that $\zeta_A(s_2, \dots, s_r)$ must be Eulerian (see Corollary 4.4.3). This fact was originally conjectured by Lara Rodríguez and Thakur [LRT14].

In [C14], the first author defined Carlitz multiple polylogarithms (abbreviated CMPL's) that are generalizations of Carlitz polylogarithms studied in [AT90]. Unlike the classical case, where there is a simple identity between multiple zeta values and multiple polylogarithms at $(1, \dots, 1)$, the function field situation is more subtle. Anderson and Thakur [AT90] showed that each Carlitz zeta value (itself a multizeta value of depth one)

is a k -linear combination of Carlitz polylogarithms at integral points, and it is generalized in [C14] that MZV's of arbitrary depth are k -linear combinations of Carlitz multiple polylogarithms at integral points. Following the terminology of Eulerian multizeta values, we call a nonzero value of a CMPL at an algebraic point *Eulerian* if it is a k -multiple $\tilde{\pi}$ raised to its weight power (see §4.3). In Theorem 4.3.2, we give a criterion to determine which CMPL's at algebraic points are Eulerian.

The main idea of this work comes from the perspective of t -motives. To handle the k -linear relations among the MZV's which interest us, we manage to lift these relations in a t -motivic way to $\bar{k}(t)$ -linear relations among specific power series in t (where \bar{k} is a fixed algebraic closure of k), which can be viewed as simplified analogue of the motivic MZV's in the classical theory. The key tool we use to accomplish the process is the linear independence criterion of [ABP04, Thm. 3.1.1] (the "ABP-criterion") that has been used successfully in the last decade for dealing with transcendence/algebraic independence questions in positive characteristic. The very fact that our motivic MZV's satisfy Frobenius (Galois) difference equations (by work of Anderson-Thakur [AT09]) also enables us to prove that the common denominator of the coefficients of the lifted relations is in $\mathbb{F}_q[t]$. This denominator gives rise to linear relations for the corresponding algebraic points under the $\mathbb{F}_q[t]$ -action, and we exploit this phenomenon as much as we can in §§2–3.

The paper is organized as follows. In §2, we first set up the necessary preliminaries and state the criterion, Theorem 2.5.2, which equates the $\mathbb{F}_q(\theta)$ -linear dependence of values of certain special series at $t = \theta$ to $\mathbb{F}_q[t]$ -linear dependence of elements of certain Ext^1 -modules. We apply [ABP04, Thm. 3.1.1] to give a proof of Theorem 2.5.2 in §3. We then apply Theorem 2.5.2 in §4 to establish the criteria for Eulerian MZV's, CMPL's at algebraic points to be Eulerian and zeta-like MZV's. Passing to t -modules in §5 we reformulate these criteria. In §6, we further prove that our criterion for Eulerian MZV's yields an algorithm for determining whether any given MZV is Eulerian or non-Eulerian. A rule specifying all Eulerian multizeta values is drawn from the data collected using this algorithm.

2. PRELIMINARIES AND STATEMENT OF THE MAIN RESULT

2.1. **Notation.** We adopt the following notation.

\mathbb{F}_q	= the finite field with q elements, for q a power of a prime number p .
θ, t	= independent variables.
A	= $\mathbb{F}_q[\theta]$, the polynomial ring in the variable θ over \mathbb{F}_q .
A_+	= set of monic polynomials in A .
k	= $\mathbb{F}_q(\theta)$, the fraction field of A .
k_∞	= $\mathbb{F}_q((1/\theta))$, the completion of k with respect to the place at infinity.
\bar{k}_∞	= a fixed algebraic closure of k_∞ .
\bar{k}	= the algebraic closure of k in \bar{k}_∞ .
\mathbb{C}_∞	= the completion of \bar{k}_∞ with respect to the canonical extension of ∞ .
$ \cdot _\infty$	= a fixed absolute value for the completed field \mathbb{C}_∞ so that $ \theta _\infty = q$.
deg	= function assigning to $x \in k_\infty$ its degree in θ .
$\mathbb{C}_\infty[[t]]$	= ring of formal power series in t over \mathbb{C}_∞ .
$\mathbb{C}_\infty((t))$	= field of Laurent series in t over \mathbb{C}_∞ .

We consider the following characteristic p multizeta values defined by Thakur [To4]: for any r -tuple of positive integers $(s_1, \dots, s_r) \in \mathbb{N}^r$,

$$(2.1.1) \quad \zeta_A(s_1, \dots, s_r) := \sum \frac{1}{a_1^{s_1} \cdots a_r^{s_r}} \in k_\infty,$$

where the sum is over $(a_1, \dots, a_r) \in A_+^r$ with $\deg a_1 > \cdots > \deg a_r$. Thakur [Toga] showed that each multizeta value is non-vanishing.

2.2. Frobenius modules. We consider the following automorphism of the field of Laurent series over \mathbb{C}_∞ , which is referred to as *Frobenius twisting*:

$$\begin{aligned} \mathbb{C}_\infty((t)) &\rightarrow \mathbb{C}_\infty((t)), \\ f := \sum_i a_i t^i &\mapsto f^{(-1)} := \sum_i a_i^{\frac{1}{q}} t^i. \end{aligned}$$

We extend Frobenius twisting to matrices with entries in $\mathbb{C}_\infty((t))$ by twisting entry-wise.

We let $\bar{k}[t, \sigma] = \bar{k}[t][\sigma]$ be the non-commutative $\bar{k}[t]$ -algebra generated by the new variable σ subject to the relation

$$\sigma f = f^{(-1)} \sigma, \quad f \in \bar{k}[t].$$

We call a left $\bar{k}[t, \sigma]$ -module a *Frobenius module* if it is free of finite rank over $\bar{k}[t]$. Morphisms of Frobenius modules are left $\bar{k}[t, \sigma]$ -module homomorphisms. We denote by \mathcal{F} the category of Frobenius modules.

The trivial object of \mathcal{F} is denoted by $\mathbf{1}$, where the underlying space of $\mathbf{1}$ is $\bar{k}[t]$ equipped with the σ -action given by

$$\sigma(f) := f^{(-1)}, \quad f \in \mathbf{1}.$$

Another example of an object in \mathcal{F} is the n -th tensor power of the Carlitz motive denoted by $\mathbb{C}^{\otimes n}$, where n is a positive integer. The underlying space of $\mathbb{C}^{\otimes n}$ is $\bar{k}[t]$, on which the action of σ is given by

$$\sigma(f) := (t - \theta)^n f^{(-1)}, \quad f \in \mathbb{C}^{\otimes n}.$$

In what follows, an object M in \mathcal{F} is said to be defined by a matrix $\Phi \in \text{Mat}_r(\bar{k}[t])$ if M is free of rank r over $\bar{k}[t]$ and the σ -action on a given $\bar{k}[t]$ -basis of M is represented by the matrix Φ .

For a Frobenius module M , we consider the tensor product $\bar{k}(t) \otimes_{\bar{k}[t]} M$ on which σ acts diagonally. It follows that $\bar{k}(t) \otimes_{\bar{k}[t]} M$ becomes a left $\bar{k}(t)[\sigma]$ -module, where $\bar{k}(t)[\sigma]$ is the twisted polynomial ring in σ over $\bar{k}(t)$ subject to the relation $\sigma h = h^{(-1)} \sigma$ for $h \in \bar{k}(t)$. The following proposition is a slight generalization of [Po8, Prop. 3.4.5], but it is crucial while proving Theorem 2.5.2.

Proposition 2.2.1. *For $i = 1, 2$, let M_i be a Frobenius module of rank r_i over $\bar{k}[t]$ defined by a given matrix $\Phi_i \in \text{Mat}_{r_i}(\bar{k}[t])$ with respect to a fixed $\bar{k}[t]$ -basis \mathbf{m}_i of M_i . Put $\mathcal{M}_i := \bar{k}(t) \otimes_{\bar{k}[t]} M_i$ for $i = 1, 2$, and let $f : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ be a homomorphism of left $\bar{k}(t)[\sigma]$ -modules. With respect to the bases $1 \otimes \mathbf{m}_1$ and $1 \otimes \mathbf{m}_2$, f is represented by a matrix $F \in \text{Mat}_{r_1 \times r_2}(\bar{k}[t])$. Suppose that $\det \Phi_i = c_i (t - \theta)^{s_i}$ for some $c_i \in \bar{k}^\times$ and $s_i \in \mathbb{Z}_{\geq 0}$ for $i = 1, 2$. Then the common denominator of the entries of F is in $\mathbb{F}_q[t]$.*

Proof. (cf. proof of [Po8, Prop. 3.4.5]) Note that since f is $\bar{k}(t)[\sigma]$ -linear, we have that

$$F^{(-1)}\Phi_2 = \Phi_1 F.$$

For a matrix $B \in \text{Mat}_{r \times s}(\bar{k}(t))$, we denote by $\text{den}(B)$ the monic least common multiple of the denominators of the entries of B . Since by hypothesis $\det \Phi_2 = c_2(t - \theta)^{s_2}$ for some $c_2 \in \bar{k}^\times$ and $s_2 \geq 0$, we find that

$$\text{den}(F)(t - \theta)^{s_2} F^{(-1)} = \text{den}(F)(t - \theta)^{s_2} \Phi_1 F \Phi_2^{-1} \in \text{Mat}_{r_1 \times r_2}(\bar{k}[t]).$$

It follows that $\text{den}(F^{(-1)})$ divides $\text{den}(F)(t - \theta)^{s_2}$. As we have $\text{den}(F^{(-1)}) = \text{den}(F)^{(-1)}$, it follows that $\deg_t(\text{den}(F^{(-1)})) = \deg_t(\text{den}(F)^{(-1)})$. Therefore, it suffices to show that $\text{den}(F^{(-1)})$ is relatively prime to $t - \theta$, since then $\text{den}(F^{(-1)}) = \text{den}(F)$, which implies $\text{den}(F) \in \mathbb{F}_q[t]$.

If $t - \theta$ divides $\text{den}(F^{(-1)})$, then $t - \theta^q$ divides $\text{den}(F)$. It forces that $t - \theta^q$ divides $\text{den}(\Phi_1 F)$, since otherwise $t - \theta^q$ would divide $\det \Phi_1 = c_1(t - \theta)^{s_1}$. Likewise, $t - \theta^q$ divides $\text{den}(\Phi_1 F \Phi_2^{-1}) = \text{den}(F^{(-1)})$. Repeating the same argument above shows that $\text{den}(F^{(-1)})$ is divisible by each of

$$t - \theta, t - \theta^q, t - \theta^{q^2}, \dots,$$

whence we obtain a contradiction since $\text{den}(F^{(-1)}) \in \bar{k}[t]$. \square

2.3. Frobenius modules connected to Carlitz multiple polylogarithms. Given a polynomial $Q := \sum_i a_i t^i \in \bar{k}[t]$, its Gauss norm is defined as $\|Q\|_\infty := \max_i \{|a_i|_\infty\}$. For a d -tuple $\mathfrak{s} = (s_1, \dots, s_d) \in \mathbb{N}^d$, we let $\mathfrak{Q} := (Q_1, \dots, Q_d) \in \bar{k}[t]^d$ satisfy the hypothesis that as $0 \leq i_d < \dots < i_1 \rightarrow \infty$,

$$(2.3.1) \quad \left(\|Q_1\|_\infty / |\theta|_\infty^{q s_1 / (q-1)} \right)^{q^{i_1}} \cdots \left(\|Q_d\|_\infty / |\theta|_\infty^{q s_d / (q-1)} \right)^{q^{i_d}} \rightarrow 0.$$

Throughout this paper, we fix a fundamental period $\tilde{\pi}$ of the Carlitz module \mathbf{C} (see [Go96, To4]). We put

$$\Omega(t) := (-\theta)^{\frac{-q}{q-1}} \prod_{i=1}^{\infty} \left(1 - \frac{t}{\theta^{q^i}} \right) \in \mathbf{C}_\infty[[t]],$$

where $(-\theta)^{\frac{1}{q-1}}$ is a suitable choice of $(q-1)$ -st root of $-\theta$ so that $\frac{1}{\Omega(\theta)} = \tilde{\pi}$. We note that Ω satisfies the functional equation $\Omega^{(-1)} = (t - \theta)\Omega$. Given d -tuples \mathfrak{s} and \mathfrak{Q} as above, we define the series

$$(2.3.2) \quad \begin{aligned} \mathcal{L}_{\mathfrak{s}, \mathfrak{Q}}(t) &:= \sum_{i_1 > \dots > i_d \geq 0} (\Omega^{s_d} Q_d)^{(i_d)} \cdots (\Omega^{s_1} Q_1)^{(i_1)} \\ &= \Omega^{s_1 + \dots + s_d} \sum_{i_1 > \dots > i_d \geq 0} \frac{Q_d^{(i_d)}(t) \cdots Q_1^{(i_1)}(t)}{((t - \theta^q) \cdots (t - \theta^{q^{i_d}}))^{s_d} \cdots ((t - \theta^q) \cdots (t - \theta^{q^{i_1}}))^{s_1}}. \end{aligned}$$

We define \mathcal{E} to be the ring consisting of formal power series $\sum_{n=0}^{\infty} a_n t^n \in \bar{k}[[t]]$ such that

$$\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|_\infty} = 0, \quad [k_\infty(a_0, a_1, a_2, \dots) : k_\infty] < \infty.$$

Then any f in \mathcal{E} has an infinite radius of convergence with respect to $|\cdot|_\infty$, and functions in \mathcal{E} are called entire functions. It is shown in [C14, Lem. 5.3.1] that the series

$\mathcal{L}_{\mathfrak{s}, \Omega}$ defined above is an entire function. We note that when $\Omega \in (\bar{k}^\times)^d$ satisfies (2.3.1) then $\tilde{\pi}^{s_1+\dots+s_r} \mathcal{L}_{\mathfrak{s}, \Omega}(\theta)$ is the Carlitz multiple polylogarithm $\text{Li}_{\mathfrak{s}}$ evaluated at the algebraic point Ω . See §4.3 for additional details.

Proposition 2.3.3. *Let $\mathfrak{s} \in \mathbb{N}^d$ and $\Omega \in \bar{k}[t]^d$ satisfy the hypothesis (2.3.1). Then for any non-negative integer N , we have that*

$$\mathcal{L}_{\mathfrak{s}, \Omega}(\theta^{q^N}) = \mathcal{L}_{\mathfrak{s}, \Omega}(\theta)^{q^N}.$$

Proof. The proof is essentially the same as the proof of [C14, Lem. 5.3.5] by changing u_i to Q_i . We omit the details. \square

Let r be a positive integer. We fix two r -tuples $\mathfrak{s} \in \mathbb{N}^r$ and $\Omega \in \bar{k}[t]^r$ satisfying (2.3.1). We define the matrix $\Phi = \Phi_{\mathfrak{s}, \Omega} \in \text{Mat}_{r+1}(\bar{k}[t])$,

$$(2.3.4) \quad \Phi := \begin{pmatrix} (t-\theta)^{s_1+\dots+s_r} & 0 & 0 & \cdots & 0 \\ Q_1^{(-1)}(t-\theta)^{s_1+\dots+s_r} & (t-\theta)^{s_2+\dots+s_r} & 0 & \cdots & 0 \\ 0 & Q_2^{(-1)}(t-\theta)^{s_2+\dots+s_r} & \ddots & & \vdots \\ \vdots & & \ddots & (t-\theta)^{s_r} & 0 \\ 0 & \cdots & 0 & Q_r^{(-1)}(t-\theta)^{s_r} & 1 \end{pmatrix}.$$

Define $\Phi' = \Phi'_{\mathfrak{s}, \Omega}$ to be the square matrix of size r cut from the upper left square of Φ :

$$(2.3.5) \quad \Phi' := \begin{pmatrix} (t-\theta)^{s_1+\dots+s_r} & & & & \\ Q_1^{(-1)}(t-\theta)^{s_1+\dots+s_r} & (t-\theta)^{s_2+\dots+s_r} & & & \\ & \ddots & & & \\ & & \ddots & & \\ & & & Q_{r-1}^{(-1)}(t-\theta)^{s_{r-1}+s_r} & (t-\theta)^{s_r} \end{pmatrix}.$$

In what follows, to avoid heavy notation we omit the subscripts \mathfrak{s} , Ω when it is clear from the context.

For $1 \leq \ell < j \leq r+1$, we define the series

$$(2.3.6) \quad \mathcal{L}_{j\ell}(t) := \sum_{i_\ell > \dots > i_{j-1} \geq 0} (\Omega^{s_{j-1}} Q_{j-1})^{(i_{j-1})} \cdots (\Omega^{s_\ell} Q_\ell)^{(i_\ell)} \in \mathcal{E},$$

which is the same series in (2.3.2) associated to the two tuples (s_ℓ, \dots, s_{j-1}) and (Q_ℓ, \dots, Q_{j-1}) . Define $\Psi \in \text{Mat}_r(\mathcal{E}) \cap \text{GL}_r(\mathbb{T})$ by

$$(2.3.7) \quad \Psi := \begin{pmatrix} \Omega^{s_1+\dots+s_r} & & & & \\ \Omega^{s_2+\dots+s_r} \mathcal{L}_{21} & \Omega^{s_2+\dots+s_r} & & & \\ \vdots & \Omega^{s_3+\dots+s_r} \mathcal{L}_{32} & \ddots & & \\ \vdots & \vdots & \ddots & \ddots & \\ \Omega^{s_r} \mathcal{L}_{r1} & \Omega^{s_r} \mathcal{L}_{r2} & \cdots & \cdots & \Omega^{s_r} \\ \mathcal{L}_{(r+1),1} & \mathcal{L}_{(r+1),2} & \cdots & \cdots & \mathcal{L}_{(r+1),r} & 1 \end{pmatrix},$$

and note that we have $\Psi^{(-1)} = \Phi\Psi$ (cf. [AT09, §2.5]). Let Ψ' be the square matrix of size r cut from the upper left square of Ψ . So then $\Psi'^{(-1)} = \Phi'\Psi'$. Note that Φ defines an object in \mathcal{F} which is a t -motive in the sense of [Po8].

2.4. **The Ext^1 -module.** We continue the notation from the previous paragraphs. We denote by M and M' the objects in \mathcal{F} defined by the matrices Φ and Φ' respectively. Note that M fits into the short exact sequence of Frobenius modules,

$$0 \rightarrow M' \rightarrow M \rightarrow \mathbf{1} \rightarrow 0,$$

and so M represents a class in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$. The group $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$ has a natural $\mathbb{F}_q[t]$ -module structure coming from Baer sum and the pushout of morphisms of M' . More precisely, if M_1 and M_2 represent classes in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$ and are defined by the two matrices respectively

$$\Phi_1 := \begin{pmatrix} \Phi' & 0 \\ \mathbf{v}_1 & 1 \end{pmatrix}, \quad \Phi_2 := \begin{pmatrix} \Phi' & 0 \\ \mathbf{v}_2 & 1 \end{pmatrix},$$

then the Baer sum $M_1 +_B M_2$ is the object in \mathcal{F} defined by the matrix

$$\begin{pmatrix} \Phi' & 0 \\ \mathbf{v}_1 + \mathbf{v}_2 & 1 \end{pmatrix}.$$

Furthermore, for any $a \in \mathbb{F}_q[t]$ multiplication by a induces an endomorphism of M' , and so the pushout $a * M_1 \in \mathcal{F}$, which is defined by the matrix

$$\begin{pmatrix} \Phi' & 0 \\ a\mathbf{v}_1 & 1 \end{pmatrix},$$

thus inducing a left $\mathbb{F}_q[t]$ -module structure on $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.

2.5. **The main theorem.** We continue with the notation as above, but assume that $r \geq 2$. We let $w := \sum_{i=1}^r s_i$ and let $Q \in \bar{k}[t]$ satisfy $\|Q\|_{\infty} < |\theta|_{\infty}^{wq/(q-1)}$. We further assume that the series $\mathcal{L}_{w,Q}(t) \in \mathcal{E}$ associated to w and Q is non-vanishing at $t = \theta$. We let $N \in \mathcal{F}$ be the Frobenius module that is defined by the matrix

$$(2.5.1) \quad \begin{pmatrix} \Phi' & \mathbf{0} \\ \mathbf{u}_w & 1 \end{pmatrix} \in \text{Mat}_{r+1}(\bar{k}[t]),$$

where $\mathbf{u}_w := (Q^{(-1)}(t - \theta)^w, 0, \dots, 0) \in \text{Mat}_{1 \times r}(\bar{k}[t])$. Then N represents a class in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.

The following result gives a criterion for the k -linear dependence of the specific values $\{\mathcal{L}_{\mathfrak{s},\Omega}(\theta), \mathcal{L}_{w,Q}(\theta), 1\}$, which is applied to the settings of Eulerian MZV's, Eulerian CMPL's at algebraic points, and zeta-like MZV's in §4. Its proof occupies the next section.

Theorem 2.5.2. *Let $r \geq 2$ be a positive integer. We fix two r -tuples $\mathfrak{s} \in \mathbb{N}^r$ and $\Omega \in \bar{k}[t]^r$ satisfying (2.3.1). Let M and M' be the objects in \mathcal{F} defined by the matrices Φ and Φ' , as in (2.3.4) and (2.3.5). For $1 \leq \ell < j \leq r + 1$, we let $\mathcal{L}_{j\ell}$ be defined as in (2.3.6) and suppose that it satisfies the non-vanishing hypothesis*

$$(2.5.3) \quad \mathcal{L}_{j\ell}(\theta) \neq 0.$$

We let $w := \sum_{i=1}^r s_i$ and let $Q \in \bar{k}[t]$ satisfy $\|Q\|_{\infty} < |\theta|_{\infty}^{wq/(q-1)}$ and $\mathcal{L}_{w,Q}(\theta) \neq 0$. Let $N \in \mathcal{F}$ be defined by the matrix given in (2.5.1). Then the following hold.

- (a) *The set $\{\mathcal{L}_{\mathfrak{s},\Omega}(\theta), \mathcal{L}_{w,Q}(\theta), 1\}$ is linearly dependent over k if and only if the classes of M and N are $\mathbb{F}_q[t]$ -linearly dependent in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$, i.e., there exists $a, b \in \mathbb{F}_q[t]$ (not both zero) so that $a * M +_B b * N$ represents a trivial class in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.*

(b) If $\{\mathcal{L}_{s,\Omega}(\theta), \mathcal{L}_{w,Q}(\theta), 1\}$ are linearly dependent over k , then each of $\mathcal{L}_{r+1,2}(\theta), \dots, \mathcal{L}_{r+1,r}(\theta)$ is also in k .

Remark 2.5.4. Note that the k -linear dependence of $\{\mathcal{L}_{s,\Omega}(\theta), \mathcal{L}_{w,Q}(\theta), 1\}$ is equivalent to the k -linear dependence of $\{\tilde{\pi}^w \mathcal{L}_{s,\Omega}(\theta), \tilde{\pi}^w \mathcal{L}_{w,Q}(\theta), \tilde{\pi}^w\}$. We mention that the values $\tilde{\pi}^w \mathcal{L}_{s,\Omega}(\theta), \tilde{\pi}^w \mathcal{L}_{w,Q}(\theta)$ satisfy the MZ property with weight w in the sense of [C14, Def. 3.4.1]), and hence by [C14, Prop. 4.3.1] we have that $\{\mathcal{L}_{s,\Omega}(\theta), \mathcal{L}_{w,Q}(\theta), 1\}$ are linearly dependent over k if and only if they are linearly dependent over \bar{k} , and that $\mathcal{L}_{s,\Omega}(\theta) \in k$ if and only if $\mathcal{L}_{s,\Omega}(\theta) \in \bar{k}$.

For the applications to Eulerian MZV's and Eulerian CMPL's at algebraic points we single out the following result, which is a special case of the theorem above.

Corollary 2.5.5. *Let notation and assumptions be given as in Theorem 2.5.2. Then we have that $\mathcal{L}_{s,\Omega}(\theta) (= \mathcal{L}_{r+1,1}(\theta))$ is in k if and only if M represents a torsion element in the $\mathbb{F}_q[t]$ -module $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.*

Proof. The proof of (\Rightarrow) is given in the case (II) of the proof of (\Rightarrow) of Theorem 2.5.2(a). The proof of (\Leftarrow) follows from the proof of (\Leftarrow) of Theorem 2.5.2(a) by putting $b = 0$. \square

3. PROOF OF THEOREM 2.5.2

3.1. A remark. For $a, b \in \mathbb{F}_q[t]$, the Frobenius module $a * M +_B b * N$ is defined by the matrix

$$(3.1.1) \quad X := \begin{pmatrix} \Phi' & \mathbf{0} \\ \mathbf{u} & 1 \end{pmatrix} \in \text{Mat}_{r+1}(\bar{k}[t]),$$

where $\mathbf{u} := (bQ^{(-1)}(t-\theta)^w, 0, \dots, 0, aQ_r^{(-1)}(t-\theta)^{s_r}) \in \text{Mat}_{1 \times r}(\bar{k}[t])$. It follows that the Frobenius module $a * M +_B b * N$ represents a trivial class in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$ if and only if there exists $\delta_1, \dots, \delta_r \in \bar{k}[t]$ so that

$$(3.1.2) \quad \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ \delta_1 & \cdots & \delta_r & 1 \end{pmatrix}^{(-1)} X = \begin{pmatrix} \Phi' & \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ \delta_1 & \cdots & \delta_r & 1 \end{pmatrix},$$

which is equivalent to that

$$(3.1.3) \quad \begin{aligned} \delta_1 &:= \delta_1^{(-1)}(t-\theta)^w + \delta_2^{(-1)}Q_1^{(-1)}(t-\theta)^w + bQ^{(-1)}(t-\theta)^w; \\ \delta_2 &:= \delta_2^{(-1)}(t-\theta)^{s_2+\dots+s_r} + \delta_3^{(-1)}Q_2^{(-1)}(t-\theta)^{s_2+\dots+s_r}; \\ &\vdots \\ \delta_{r-1} &:= \delta_{r-1}^{(-1)}(t-\theta)^{s_{r-1}+s_r} + \delta_r^{(-1)}Q_{r-1}^{(-1)}(t-\theta)^{s_{r-1}+s_r}; \\ \delta_r &:= \delta_r^{(-1)}(t-\theta)^{s_r} + aQ_r^{(-1)}(t-\theta)^{s_r}. \end{aligned}$$

3.2. **Proof of Theorem 2.5.2 (a)(\Rightarrow).** Suppose that $\{\mathcal{L}_{s,\Omega}(\theta), \mathcal{L}_{w,Q}(\theta), 1\}$ are linearly dependent over k . Our goal is to find $a, b \in \mathbb{F}_q[t]$ (not both zero) and $\delta_1, \dots, \delta_r \in \bar{k}[t]$ satisfying the equations (3.1.3).

Define the matrix

$$\tilde{\Phi} := \begin{pmatrix} 1 & & & & \\ & \Phi & & & \\ & Q^{(-1)}(t-\theta)^w, 0, \dots, 0 & & & \\ & & & & 1 \end{pmatrix} \in \text{Mat}_{r+3}(\bar{k}[t])$$

and put

$$\tilde{\psi} := \begin{pmatrix} 1 \\ \Omega^{s_1+\dots+s_r} \\ \Omega^{s_2+\dots+s_r} \mathcal{L}_{21} \\ \vdots \\ \mathcal{L}_{r+1,1} \\ \mathcal{L}_{w,Q} \end{pmatrix}.$$

Then we have the difference equations $\tilde{\psi}^{(-1)} = \tilde{\Phi}\tilde{\psi}$.

Note that $\mathcal{L}_{r+1,1} = \mathcal{L}_{s,\Omega}$. By [ABP04, Thm. 3.1.1] there exists

$$\mathbf{f} = (f_0, f_1, \dots, f_{r+2}) \in \text{Mat}_{1 \times (r+3)}(\bar{k}[t])$$

so that $\mathbf{f}\tilde{\psi} = 0$ and $\mathbf{f}(\theta)\tilde{\psi}(\theta) = 0$, which describes a nontrivial k -linear relation among $\{\mathcal{L}_{s,\Omega}(\theta), \mathcal{L}_{w,Q}(\theta), 1\}$. Note that

$$f_1(\theta) = \dots = f_r(\theta) = 0.$$

Now we assume that $f_{r+2}(\theta) \neq 0$. Note if $f_{r+2}(\theta) = 0$, then we have that

$$\{\mathcal{L}_{s,\Omega}(\theta) = \mathcal{L}_{r+1,1}(\theta), 1\}$$

are linearly dependent over k , i.e., $\mathcal{L}_{r+1,1}(\theta) \in k$, and this case will be included in the case (II) below.

If we put $\tilde{\mathbf{f}} := \frac{1}{f_{r+2}}\mathbf{f} \in \text{Mat}_{1 \times (r+3)}(\bar{k}(t))$, then all entries of $\tilde{\mathbf{f}}$ are regular at $t = \theta$. Considering the Frobenius twisting-action $(\cdot)^{(-1)}$ on the equation $\tilde{\mathbf{f}}\tilde{\psi} = 0$ and subtracting the resulting equation from $\tilde{\mathbf{f}}\tilde{\psi} = 0$, we obtain

$$(3.2.1) \quad (\tilde{\mathbf{f}} - \tilde{\mathbf{f}}^{(-1)}\tilde{\Phi})\tilde{\psi} = 0.$$

Explicit calculations show that $\tilde{\mathbf{f}} - \tilde{\mathbf{f}}^{(-1)}\tilde{\Phi} = (B, B_1, \dots, B_{r+1}, 0)$, where

$$(3.2.2) \quad \begin{aligned} B &:= \frac{f_0}{f_{r+2}} - \left(\frac{f_0}{f_{r+2}}\right)^{(-1)} \\ B_1 &:= \frac{f_1}{f_{r+2}} - \left(\frac{f_1}{f_{r+2}}\right)^{(-1)}(t-\theta)^w - \left(\frac{f_2}{f_{r+2}}\right)^{(-1)}Q_1^{(-1)}(t-\theta)^w - Q^{(-1)}(t-\theta)^w; \\ B_2 &:= \frac{f_2}{f_{r+2}} - \left(\frac{f_2}{f_{r+2}}\right)^{(-1)}(t-\theta)^{s_2+\dots+s_r} - \left(\frac{f_3}{f_{r+2}}\right)^{(-1)}Q_2^{(-1)}(t-\theta)^{s_2+\dots+s_r}; \\ &\vdots \\ B_r &:= \frac{f_r}{f_{r+2}} - \left(\frac{f_r}{f_{r+2}}\right)^{(-1)}(t-\theta)^{s_r} - \left(\frac{f_{r+1}}{f_{r+2}}\right)^{(-1)}Q_r^{(-1)}(t-\theta)^{s_r} \\ B_{r+1} &:= \left(\frac{f_{r+1}}{f_{r+2}}\right) - \left(\frac{f_{r+1}}{f_{r+2}}\right)^{(-1)}. \end{aligned}$$

Case (I). $\mathcal{L}_{r+1,1}(\theta) \notin k$ (which is equivalent to $\mathcal{L}_{r+1,1}(\theta) \notin \bar{k}$ by Remark 2.5.4). We claim that $B = B_1 = \cdots = B_{r+1} = 0$. Assuming this claim first, we see that

$$(3.2.3) \quad \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ f_0/f_{r+2} & \cdots & f_{r+1}/f_{r+2} & 1 \end{pmatrix}^{(-1)} \tilde{\Phi} = \begin{pmatrix} 1 & & \\ & \Phi & \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ f_0/f_{r+2} & \cdots & f_{r+1}/f_{r+2} & 1 \end{pmatrix},$$

where Φ' is defined in (2.3.5). Let \tilde{M} be the Frobenius module defined by the matrix $\tilde{\Phi}$. Then the equation (3.2.3) gives a left $\bar{k}(t)[\sigma]$ -module homomorphism between $\bar{k}(t) \otimes_{\bar{k}[t]} (\mathbf{1} \oplus M \oplus \mathbf{1})$ and $\bar{k}(t) \otimes_{\bar{k}[t]} \tilde{M}$. It follows that by Proposition 2.2.1 we have that the denominator of each f_i/f_{r+2} is in $\mathbb{F}_q[t]$ for $i = 0, \dots, r+1$. Now we let $b \in \mathbb{F}_q[t]$ be the common denominator of $f_0/f_{r+2}, \dots, f_{r+1}/f_{r+2}$, and take $\delta_i := bf_i/f_{r+2} \in \bar{k}[t]$ for $i = 1, \dots, r$. Note that the vanishing of B_{r+1} implies $f_{r+1}/f_{r+2} \in \mathbb{F}_q(t)$ and hence $a := bf_{r+1}/f_{r+2} \in \mathbb{F}_q[t]$. Multiplying by b on the both sides of (3.2.2) one finds exactly the identities (3.1.3), which imply that $a * M +_B b * N$ represents a trivial class in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.

To prove the claim above, we consider (3.2.1), which is expanded as

$$(3.2.4) \quad B + B_1 \Omega^{s_1 + \cdots + s_r} + B_2 \Omega^{s_2 + \cdots + s_r} \mathcal{L}_{21} + \cdots + B_r \Omega^{s_r} \mathcal{L}_{r1} + B_{r+1} \mathcal{L}_{r+1,1} = 0.$$

For each $1 \leq \ell < j \leq r+1$ and any non-negative integer n , by Proposition 2.3.3

$$(3.2.5) \quad \mathcal{L}_{j\ell}(\theta^{q^n}) = \mathcal{L}_{j\ell}(\theta)^{q^n},$$

which is nonzero by hypothesis. Since B and each B_i are rational functions in $\bar{k}(t)$, B and B_i are defined at $t = \theta^{q^n}$ for sufficiently large integers n . We further note that Ω has a simple zero at $t = \theta^{q^i}$ for each positive integer i . Specializing (3.2.4) at $t = \theta^{q^n}$ shows that $B(\theta^{q^n}) = B_{r+1}(\theta^{q^n}) = 0$ for any $n \gg 0$ because of $\mathcal{L}_{r+1,1}(\theta) \notin \bar{k}$. This follows that $B = B_{r+1} = 0$.

Next, dividing (3.2.4) by Ω^{s_r} and then specializing at $t = \theta^{q^n}$, we see from (3.2.5) that $B_r(\theta^{q^n}) = 0$ for all sufficiently large integers n . It follows that $B_r = 0$. Furthermore, using (3.2.4) and repeating the arguments above we can show that $B_{r-1} = \cdots = B_1 = 0$, whence the desired claim.

Case (II). $\mathcal{L}_{r+1,1}(\theta) \in k$. In this case, we apply [ABP04, Thm. 3.1.1] to the difference equations

$$\begin{pmatrix} 1 \\ \Omega^{s_1 + \cdots + s_r} \\ \Omega^{s_2 + \cdots + s_r} \mathcal{L}_{21} \\ \vdots \\ \mathcal{L}_{r+1,1} \end{pmatrix}^{(-1)} = \begin{pmatrix} 1 & \\ & \Phi \end{pmatrix} \begin{pmatrix} 1 \\ \Omega^{s_1 + \cdots + s_r} \\ \Omega^{s_2 + \cdots + s_r} \mathcal{L}_{21} \\ \vdots \\ \mathcal{L}_{r+1,1} \end{pmatrix}.$$

The arguments are the similar in this case as in the previous one, and we omit them. Moreover, they exactly show that the class of M is a torsion element in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.

3.3. Proof of Theorem 2.5.2 (a)(\Leftrightarrow) and (b). Suppose that there exist $a, b \in \mathbb{F}_q[t]$ (not both zero) for which $a * M +_B b * N$ represents a trivial class in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$. Note that

Since we are working in characteristic p , taking the q^n -th root of both sides of (3.3.2) shows that $\mathcal{L}_{r+1,1}(\theta)(= \mathcal{L}_{s,\Omega}(\theta))$ and $\mathcal{L}_{w,Q}(\theta)$ are linearly dependent over k , and each of $\mathcal{L}_{r+1,2}(\theta), \dots, \mathcal{L}_{r+1,r}(\theta)$ is rational in k .

4. APPLICATIONS TO MULTIZETA VALUES AND MULTIPLE POLYLOGARITHMS

In this section, we apply Theorem 2.5.2 to establish criteria for MZV's and CMPL's at algebraic points to be Eulerian and for MZV's to be zeta-like.

4.1. Anderson-Thakur polynomials. Define $D_0 = 1$ and $D_i := \prod_{j=0}^{i-1} (\theta^{q^i} - \theta^{q^j})$ for $i \in \mathbb{N}$. For a non-negative integer n , we express n as

$$n = \sum_{i=0}^{\infty} n_i q^i \quad (0 \leq n_i \leq q-1, n_i = 0 \text{ for } i \gg 0),$$

and we recall the definition of the arithmetic Γ -function,

$$\Gamma_{n+1} := \prod_{i=0}^{\infty} D_i^{n_i} \in A.$$

We put $G_0(y) := 1$ and define polynomials $G_n(y) \in \mathbb{F}_q[t, y]$ for $n \in \mathbb{N}$ by the product

$$G_n(y) = \prod_{i=1}^n (t^{q^n} - y^{q^i}).$$

Note that $G_{n+1}(y^q) = (t - y^q)^{q^{n+1}} G_n(y)^q$.

For $n = 0, 1, 2, \dots$ we define the sequence of Anderson-Thakur polynomials $H_n \in A[t]$ by the generating function identity

$$\left(1 - \sum_{i=0}^{\infty} \frac{G_i(\theta)}{D_i|_{\theta=t}} x^{q^i} \right)^{-1} = \sum_{n=0}^{\infty} \frac{H_n}{\Gamma_{n+1}|_{\theta=t}} x^n.$$

We note that for $0 \leq n \leq q-1$ we have $H_n = 1$. We have made a change of notation by $t \leftarrow T, \theta \leftarrow y$ from [AT90, (3.7.1)] in order to match better the notation in [AT09], [C14]. It is shown in [AT90, 3.7.3] that when one regards H_n as a polynomial in θ over $\mathbb{F}_q[t]$, one has $\deg_{\theta} H_n \leq \frac{nq}{q-1}$, whence

$$(4.1.1) \quad \|H_n\|_{\infty} \leq |\theta|_{\infty}^{nq/(q-1)}.$$

The crucial identity developed in [AT90], [AT09] is that

$$(H_{s-1}\Omega^s)^{(d)}(\theta) = \frac{\Gamma_s S_d(s)}{\tilde{\pi}^s}, \quad \forall s, d \in \mathbb{N},$$

where $S_d(s)$ is the power sum

$$S_d(s) := \sum_{\substack{a \in A_+ \\ \deg a = d}} \frac{1}{a^s} \in k.$$

It follows that if we put $\Omega = (H_{s_1-1}, \dots, H_{s_r-1})$, then by (4.1.1) Ω satisfies the hypothesis (2.3.1). Furthermore, specialization of the series

$$\mathcal{L}_{s,\Omega} := \sum_{i_1 > \dots > i_r \geq 0} (\Omega^{s_r} H_{s_r-1})^{(i_r)} \dots (\Omega^{s_1} H_{s_1-1})^{(i_1)}$$

at $t = \theta$ is equal to

$$\Gamma_{s_1} \cdots \Gamma_{s_r} \zeta_A(s_1, \dots, s_r) / \tilde{\pi}^{s_1 + \cdots + s_r}.$$

4.2. A criterion for Eulerian MZV's. We continue with the notation defined in the previous section. Given $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$, we let $\Omega = (H_{s_1-1}, \dots, H_{s_r-1})$ and M (resp. M') be defined by Φ as in (2.3.4) (resp. by Φ' as in (2.3.5)). For this choice of Ω each series $\mathcal{L}_{j\ell}$ defined in (2.3.6) evaluates at $t = \theta$ as

$$\mathcal{L}_{j\ell}(\theta) = \Gamma_{s_\ell} \cdots \Gamma_{s_{j-1}} \zeta_A(s_\ell, \dots, s_{j-1}) / \tilde{\pi}^{s_\ell + \cdots + s_{j-1}}.$$

These values are non-vanishing by the work of Thakur [Toga, Thm. 4], and so it satisfies the non-vanishing hypothesis (2.5.3). In particular,

$$(4.2.1) \quad \begin{aligned} \mathcal{L}_{r+1,1}(\theta) &= \Gamma_{s_1} \cdots \Gamma_{s_r} \zeta_A(s_1, \dots, s_r) / \tilde{\pi}^{s_1 + \cdots + s_r} \\ \mathcal{L}_{r+1,2}(\theta) &= \Gamma_{s_2} \cdots \Gamma_{s_r} \zeta_A(s_2, \dots, s_r) / \tilde{\pi}^{s_2 + \cdots + s_r} \\ &\vdots \\ \mathcal{L}_{r+1,r}(\theta) &= \Gamma_{s_r} \zeta_A(s_r) / \tilde{\pi}^{s_r}. \end{aligned}$$

Note that the hypotheses in Theorem 2.5.2 are satisfied and thus applying Corollary 2.5.5 in this situation, we obtain the following criterion to determine when a given multizeta value is Eulerian.

Theorem 4.2.2. *The multizeta value $\zeta_A(s_1, \dots, s_r)$ is Eulerian if and only if the class of M is a torsion element in the $\mathbb{F}_q[t]$ -module $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.*

Applying Theorem 2.5.2(b), we obtain additional information about when a given multizeta value is Eulerian.

Corollary 4.2.3. *Suppose that $\zeta_A(s_1, \dots, s_r)$ is Eulerian. Then the following hold.*

(a) *Each of the multizeta values*

$$\zeta_A(s_2, \dots, s_r), \zeta_A(s_3, \dots, s_r), \dots, \zeta_A(s_r)$$

is also Eulerian.

(b) *Each $\zeta_A(s_i)$ is also Eulerian for $i = 1, \dots, r$. That is, each s_i is divisible by $q - 1$.*

Proof. Using (4.2.1), part (a) follows from Theorem 2.5.2(b). For part (b), we note that $\tilde{\pi}^n \in k_\infty$ if and only if $(q - 1) \mid n$. If $\zeta_A(s_1, \dots, s_r)$ is Eulerian, then part (a) implies that so are $\zeta_A(s_2, \dots, s_r), \zeta_A(s_3, \dots, s_r), \dots, \zeta_A(s_r)$. Therefore, we have the divisibility relations

$$(q - 1) \mid (s_1 + \cdots + s_r), \dots, (q - 1) \mid (s_{r-1} + s_r), (q - 1) \mid s_r.$$

It follows that each s_i is divisible by $q - 1$ for $i = 1, \dots, r$, and so each $\zeta_A(s_i)$ is Eulerian by the original results of Carlitz [Ca35]. \square

4.3. Application to multiple polylogarithms at algebraic points. We now consider Carlitz multiple polylogarithms [C14], which are generalizations of Carlitz polylogarithms [Ca35, AT90]. We state a criterion to determine when a nonzero Carlitz multiple polylogarithm at an algebraic point is Eulerian and show how to deduce it from Corollary 2.5.5.

Define $L_0 := 1$ and $L_i := (\theta - \theta^q) \cdots (\theta - \theta^{q^i})$ for $i \in \mathbb{N}$. Given an r -tuple of positive integers $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$, its associated Carlitz multiple polylogarithm, abbreviated CMPL, is defined by

$$(4.3.1) \quad \text{Li}_{\mathfrak{s}}(z_1, \dots, z_r) := \sum_{i_1 > \dots > i_r \geq 0} \frac{z_1^{q^{i_1}} \cdots z_r^{q^{i_r}}}{L_{i_1}^{s_1} \cdots L_{i_r}^{s_r}}.$$

We further denote by

$$\mathbb{D}_{\mathfrak{s}} := \{\mathbf{u} = (u_1, \dots, u_r) \in \mathbb{C}_{\infty}^r \mid \text{Li}_{\mathfrak{s}}(\mathbf{u}) \text{ converges}\},$$

the convergence domain of $\text{Li}_{\mathfrak{s}}$. We can describe $\mathbb{D}_{\mathfrak{s}}$ as

$$\mathbb{D}_{\mathfrak{s}} = \left\{ \mathbf{u} \in \mathbb{C}_{\infty}^r \mid |u_1 / (\theta^{q^{s_1}})|_{\infty}^{q^{i_1}} \cdots |u_r / (\theta^{q^{s_r}})|_{\infty}^{q^{i_r}} \rightarrow 0 \text{ as } 0 \leq i_r < \dots < i_1 \rightarrow \infty \right\}.$$

The weight of $\text{Li}_{\mathfrak{s}}$ is defined to be $\text{wt}(\mathfrak{s}) := \sum_{i=1}^r s_i$ and its depth is defined to be r . We do not know whether $\text{Li}_{\mathfrak{s}}(\mathbf{u})$ is non-vanishing for arbitrary $\mathbf{u} \in \mathbb{D}_{\mathfrak{s}}$, but it is non-vanishing when \mathbf{u} lies in the smaller domain

$$\mathbb{D}'_{\mathfrak{s}} := \left\{ (u_1, \dots, u_r) \in \mathbb{C}_{\infty}^r \mid |u_i|_{\infty} < q^{\frac{s_i q}{q-1}} \text{ for } i = 1, \dots, r \right\}$$

(see [C14, Remark 5.1.4]).

As a generalization of the work of Anderson and Thakur [AT90], the first author of the present paper showed that for any $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$, $\zeta_A(s_1, \dots, s_r)$ is a k -linear combination of $\text{Li}_{\mathfrak{s}}$ at some integral points in $A^r \cap \mathbb{D}_{\mathfrak{s}}$ [C14, Thm. 5.5.2]. From now on, we fix an r -tuple $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$ and $\mathbf{u} = (u_1, \dots, u_r) \in (\bar{k}^{\times})^r \cap \mathbb{D}_{\mathfrak{s}}$, and suppose that $\text{Li}_{\mathfrak{s}}(\mathbf{u}) \neq 0$. Following the terminology for multizeta values, we shall call $\text{Li}_{\mathfrak{s}}(\mathbf{u})$ *Eulerian* if the ratio $\text{Li}_{\mathfrak{s}}(\mathbf{u}) / \tilde{\pi}^{\text{wt}(\mathfrak{s})}$ lies in k . Note that it is shown in [C14] that $\text{Li}_{\mathfrak{s}}(\mathbf{u}) / \tilde{\pi}^{\text{wt}(\mathfrak{s})} \in \bar{k}$ if and only if $\text{Li}_{\mathfrak{s}}(\mathbf{u}) / \tilde{\pi}^{\text{wt}(\mathfrak{s})} \in k$.

We put $\Omega := \mathbf{u} = (u_1, \dots, u_r)$ and let M (resp. M') be the Frobenius module defined by Φ as in (2.3.4) (resp. Φ' as in (2.3.5)). For $1 \leq i < j \leq r$, we put

$$\text{Li}_{\mathfrak{s}_{ij}}(\mathbf{u}_{ij}) := \text{Li}_{(s_i, \dots, s_j)}(u_i, \dots, u_j).$$

Note that in this setting we have $\mathcal{L}_{\mathfrak{s}, \Omega}(\theta) = \text{Li}_{\mathfrak{s}}(\mathbf{u}) / \tilde{\pi}^{\text{wt}(\mathfrak{s})}$, and hence $\text{Li}_{\mathfrak{s}}(\mathbf{u})$ is Eulerian if and only if $\mathcal{L}_{\mathfrak{s}, \Omega}(\theta) \in k$. Applying Corollary 2.5.5 and Theorem 2.5.2(b), we obtain the following result.

Theorem 4.3.2. *For any $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$ and $\mathbf{u} = (u_1, \dots, u_r) \in (\bar{k}^{\times})^r \cap \mathbb{D}_{\mathfrak{s}}$, we let M and M' be defined as above. Suppose that $\text{Li}_{\mathfrak{s}_{ij}}(\mathbf{u}_{ij})$ is nonzero for all $1 \leq i < j \leq r$. Then we have*

- (a) $\text{Li}_{\mathfrak{s}}(\mathbf{u})$ is Eulerian if and only if M represents a torsion element in the $\mathbb{F}_q[t]$ -module $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.
- (b) If $\text{Li}_{\mathfrak{s}}(\mathbf{u})$ is Eulerian, then each of $\text{Li}_{(s_2, \dots, s_r)}(u_2, \dots, u_r), \dots, \text{Li}_{s_r}(u_r)$ is also Eulerian.

Remark 4.3.3. The non-vanishing hypothesis of $\text{Li}_{\mathfrak{s}_{ij}}(\mathbf{u}_{ij})$ is non-empty. Indeed, for those points $\mathbf{u} \in \mathbb{D}'_{\mathfrak{s}}$ we have that $\text{Li}_{\mathfrak{s}_{ij}}(\mathbf{u}_{ij}) \neq 0$ for all $1 \leq i < j \leq r$. See [C14, Rem. 5.1.5].

4.4. Applications to zeta-like multizeta values. In this section we apply Corollary 2.5.5 to confirm a conjecture of Lara Rodríguez and Thakur. As defined by Thakur a multizeta value $\zeta_A(s_1, \dots, s_r)$ with weight $w := \sum_{i=1}^n s_i$ is called *zeta-like* if the ratio $\zeta_A(s_1, \dots, s_r)/\zeta_A(w)$ is algebraic over k . Note that if $\zeta_A(s_1, \dots, s_r)$ is zeta-like, then by [C14, Cor. 2.3.3] the ratio $\zeta_A(s_1, \dots, s_r)/\zeta_A(w)$ is actually in k .

4.4.1. Zeta-like MZV's. For a multizeta value $\zeta_A(s_1, \dots, s_r)$, Lara Rodríguez and Thakur [LRT14] conjectured the following assertion.

Conjecture 4.4.1. *If $\zeta_A(s_1, \dots, s_r)$ is zeta-like, then $\zeta_A(s_2, \dots, s_r)$ is Eulerian.*

In what follows, we confirm this conjecture stated as Corollary 4.4.3, which is a consequence of the following theorem. For more conjectures concerning zeta-like MZV's, we refer the reader to [LRT14].

4.4.2. The setting and results. Given $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$, put $\Omega = (H_{s_1-1}, \dots, H_{s_r-1})$ and $Q := H_{w-1}$, where $\{H_n\}$ be the Anderson-Thakur polynomials given in §4.1 and $w := \sum_{i=1}^n s_i$. Note that

$$\mathcal{L}_{w,Q}(\theta) = \frac{\Gamma_w \zeta_A(w)}{\tilde{\pi}^w},$$

and so by (4.1.1) the conditions of Theorem 2.5.2 are satisfied.

Now let Φ (resp. X) be defined as (2.3.4) (resp. (3.1.1)), and let M (resp. N) be the object in \mathcal{F} defined by Φ (resp. X). Then applying Theorem 2.5.2 we have the following criterion.

Theorem 4.4.2. *Let $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$ with $r \geq 2$, and let notation be given as above. Then we have that $\{\zeta_A(s_1, \dots, s_r), \zeta_A(w), \tilde{\pi}^w\}$ are linearly dependent over k if and only if the classes of M and N are $\mathbb{F}_q[t]$ -linearly dependent in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$.*

Corollary 4.4.3. *If $\zeta_A(s_1, \dots, s_r)$ is zeta-like, then each of*

$$\zeta_A(s_2, \dots, s_r), \zeta_A(s_3, \dots, s_r) \dots, \zeta_A(s_r)$$

is also Eulerian. In particular, each of s_2, \dots, s_r is divisible by $q - 1$.

Proof. Note that

$$\mathcal{L}_{\mathfrak{s},\Omega}(\theta) = \Gamma_{s_1} \cdots \Gamma_{s_r} \zeta_A(s_1, \dots, s_r) / \tilde{\pi}^w, \quad \mathcal{L}_{w,Q}(\theta) = \Gamma_w \zeta_A(w) / \tilde{\pi}^w$$

and for each $2 \leq \ell \leq r$,

$$\mathcal{L}_{r+1,\ell}(\theta) = \Gamma_{s_\ell} \cdots \Gamma_{s_r} \zeta_A(s_\ell, \dots, s_r) / \tilde{\pi}^{s_\ell + \dots + s_r}.$$

Since by hypothesis $\zeta_A(s_1, \dots, s_r)$ is zeta-like, the result follows from Theorem 2.5.2(b). \square

5. OPERATING ON t -MODULES

The purpose of this section is to reformulate our criteria via t -modules.

5.1. The structure of rational torsion points of $C^{\otimes n}$.

5.1.1. *Definition of t -modules.* We first review the definition of t -modules [A86]. Let $\tau = (x \mapsto x^q) : \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$ be the Frobenius q -th power operator and let $\mathbb{C}_\infty[\tau]$ be the twisted polynomial ring in τ over \mathbb{C}_∞ subject to the relation $\tau\alpha = \alpha^q\tau$ for $\alpha \in \mathbb{C}_\infty$. For a positive integer d , a d -dimensional t -module is a pair (E, ϕ) , where E is the d -dimensional algebraic group \mathbb{G}_a^d and ϕ is an \mathbb{F}_q -linear ring homomorphism

$$\phi : \mathbb{F}_q[t] \rightarrow \text{Mat}_d(\mathbb{C}_\infty[\tau])$$

so that when we write $\phi_t = \alpha_0 + \sum_i \alpha_i \tau^i$ with $\alpha_i \in \text{Mat}_d(\mathbb{C}_\infty)$, $\alpha_0 - \theta I_d$ is a nilpotent matrix. In this way, $E(\mathbb{C}_\infty)$ is equipped with an $\mathbb{F}_q[t]$ -module structure via the map ϕ . For a subring $R \subset \mathbb{C}_\infty$ containing A , we say that the t -module E is defined over R if α_i lies in $\text{Mat}_d(A)$ for all $i \geq 0$.

For any d -dimensional t -module (E, ϕ) , Anderson [A86] showed that one has the \mathbb{F}_q -linear function $\exp_E : \mathbb{C}_\infty^d \rightarrow \mathbb{C}_\infty^d$ satisfying that for $\mathbf{z} = (z_1, \dots, z_d)^{\text{tr}}$ and any $a \in \mathbb{F}_q[t]$,

- $\exp_E(\mathbf{z}) \equiv \mathbf{z} \pmod{\text{degree } q}$
- $\exp_E(\partial\phi_a(\mathbf{z})) = \phi_a(\exp_E(\mathbf{z}))$,

where $\partial\phi_a$ is the differential of the morphism ϕ_a at the identity element of E . If \exp_E is surjective, then E is called uniformizable.

5.1.2. *Anderson-Thakur special points.* For a positive integer n , the n -th tensor power of the Carlitz $\mathbb{F}_q[t]$ -module denoted by $\mathbf{C}^{\otimes n}$ is an n -dimensional t -module defined over A together with the \mathbb{F}_q -linear ring homomorphism

$$[\cdot]_n : \mathbb{F}_q[t] \rightarrow \text{Mat}_n(\mathbb{C}_\infty[\tau])$$

given by

$$[t]_n = \theta I_n + N_n + E_n \tau,$$

where

$$N_n := \begin{pmatrix} 0 & 1 & \cdots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & 1 \\ 0 & \cdots & \cdots & 0 \end{pmatrix}, \quad E_n := \begin{pmatrix} 0 & \cdots & \cdots & 0 \\ \vdots & & & \vdots \\ \vdots & & & \vdots \\ 1 & \cdots & \cdots & 0 \end{pmatrix}.$$

Note that for $n = 1$, the definition above is the Carlitz $\mathbb{F}_q[t]$ -module \mathbf{C} . It is shown in [AT90, Cor. 2.5.8] that $\mathbf{C}^{\otimes n}$ is uniformizable and the kernel of $\exp_{\mathbf{C}^{\otimes n}}$ is a rank one $\mathbb{F}_q[t]$ -module (via the $\partial[a]_n$ -action) with a generator of the form

$$\lambda_n = \begin{pmatrix} * \\ \vdots \\ \tilde{\pi}^n \end{pmatrix} \in \mathbb{C}_\infty^n.$$

To connect with $\zeta_A(n)$, Anderson and Thakur defined the following special points (see [AT90, (3.8.2)]).

Definition 5.1.1. For each positive integer n , we let $H_{n-1} \in A[t]$ be the Anderson-Thakur polynomial in §4.1. We write $H_{n-1} = \sum_{i \geq 0} h_{ni} \theta^i$ with $h_{ni} \in \mathbb{F}_q[t]$. Then we define

$$Z_n := \sum_{i \geq 0} [h_{ni}]_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ \theta^i \end{pmatrix} \in \mathbf{C}^{\otimes n}(A)$$

and call it an Anderson-Thakur special point.

It is shown in [AT90, Thm. 3.8.3] there exists a vector of the form

$$z_n = \begin{pmatrix} * \\ \vdots \\ * \\ \Gamma_n \zeta_A(n) \end{pmatrix}$$

so that

$$\exp_{\mathbf{C}^{\otimes n}}(z_n) = Z_n.$$

Remark 5.1.2. For a positive integer n even, we put

$$a := \frac{\Gamma_{n+1}}{\Gamma_n} \text{den BC}(n)|_{\theta=t},$$

where $\text{den BC}(n)$ denotes the denominator of the n -th Bernoulli-Carlitz number. Then by the formula (1.0.1) of Carlitz, the property that $\text{Ker exp}_{\mathbf{C}^{\otimes n}}$ is a rank one $\mathbb{F}_q[t]$ -module, the functional equation $\exp_{\mathbf{C}^{\otimes n}}(\partial[a]_n \mathbf{z}) = [a]_n (\exp_{\mathbf{C}^{\otimes n}}(\mathbf{z}))$ and [Yu91, Thm. 2.3], we see that Z_n is an a -torsion point in $\mathbf{C}^{\otimes n}(A)$.

5.1.3. *The structure of $\mathbf{C}^{\otimes n}(k)_{\text{tor}}$.* For any nonzero polynomial $f \in \mathbb{F}_q[t]$, denote by $\mathbf{C}^{\otimes n}[f]$ the set of f -torsion elements:

$$\mathbf{C}^{\otimes n}[f] := \left\{ \mathbf{x} \in \mathbf{C}^{\otimes n}(\bar{k}) = \bar{k}^n \mid [f]_n(\mathbf{x}) = \mathbf{0} \right\}.$$

We further define the set of rational torsion points of $\mathbf{C}^{\otimes n}$:

$$\mathbf{C}^{\otimes n}(k)_{\text{tor}} := \left\{ \mathbf{x} \in \mathbf{C}^{\otimes n}(k) = k^n \mid [f]_n(\mathbf{x}) = \mathbf{0} \text{ for some nonzero } f \in \mathbb{F}_q[t] \right\}.$$

Note that in [AT90, Prop. 1.11.2] Anderson and Thakur showed that $\mathbf{C}^{\otimes n}(k)_{\text{tor}}$ is trivial if n is odd (ie., $(q-1) \nmid n$). The following result is the structure of $\mathbf{C}^{\otimes n}(k)_{\text{tor}}$ when n is even, and we thank Y.-L. Kuan for providing us a proof of the following lemma.

Lemma 5.1.3. *Let n be a positive integer divisible by $q-1$. We decompose $n = p^\ell n_1 (q^h - 1)$ where $p \nmid n_1$ and h is the greatest integer such that $(q^h - 1) \mid n$. Then*

$$\mathbf{C}^{\otimes n}(k)_{\text{tor}} = \prod_{\deg P \mid h} \mathbf{C}^{\otimes n}[P^{p^\ell}],$$

where the product runs through all monic irreducible polynomials $P \in \mathbb{F}_q[t]$ with $\deg P \mid h$. In particular, the Fitting ideal of the finite $\mathbb{F}_q[t]$ -module $\mathbf{C}^{\otimes n}(k)_{\text{tor}}$ is generated by $(t^{q^h} - t)^{p^\ell}$.

Proof. For any nonzero $f \in \mathbb{F}_q[t]$, let $(\mathbb{F}_q[t]/f)^{\times n}$ be the group of the n -th powers of elements of $(\mathbb{F}_q[t]/f)^\times$. By [AT90, Prop. 1.11.1] one knows that the Galois group $\text{Gal}(k(\mathbf{C}^{\otimes n}[f])/k)$ is isomorphic to $(\mathbb{F}_q[t]/f)^{\times n}$. It follows that $\mathbf{C}^{\otimes n}[f] \subseteq \mathbf{C}^{\otimes n}(k)$ if and only if the group $(\mathbb{F}_q[t]/f)^{\times n}$ is trivial.

Note that the group $(\mathbb{F}_q[t]/P)^{\times n}$ is non-trivial if $\deg P \nmid h$. Thus, to prove the lemma it suffices to show that the group $(\mathbb{F}_q[t]/P^{p^\ell})^{\times n}$ is trivial and $(\mathbb{F}_q[t]/P^{p^{\ell+1}})^{\times n}$ is non-trivial for any monic irreducible polynomial P with $\deg P \mid h$. Suppose that $\deg P \mid h$. Let $(\mathbb{F}_q[t]/P^{p^\ell})^{(1)}$ be the kernel of the natural map from $(\mathbb{F}_q[t]/P^{p^\ell})^\times$ to $(\mathbb{F}_q[t]/P)^\times$. Then the group $(\mathbb{F}_q[t]/P^{p^\ell})^\times$ is isomorphic to the direct product of $(\mathbb{F}_q[t]/P)^\times$ and

$(\mathbb{F}_q[t]/P^{p^\ell})^{(1)}$. Since $\deg P \mid h$, the group $(\mathbb{F}_q[t]/P)^{\times n}$ is trivial. Note that every element in $(\mathbb{F}_q[t]/P^{p^\ell})^{(1)}$ can be represented by a polynomial of the form $a = 1 + bP$ with $b \in \mathbb{F}_q[t]$. Then

$$a^n = \left((1 + bP)^{p^\ell} \right)^{n_1(q^h-1)} = \left(1 + b^{p^\ell} P^{p^\ell} \right)^{n_1(q^h-1)} \equiv 1 \pmod{P^{p^\ell}},$$

and hence the group $(\mathbb{F}_q[t]/P^{p^\ell})^{\times n}$ is trivial. On the other hand, we now consider $a = 1 + P$. Since $(\mathbb{F}_q[t]/P^{p^{\ell+1}})^{(1)}$ is a p -group, $a^n \neq 1$ in $(A/P^{p^{\ell+1}})^{(1)}$ if and only if $a^{p^\ell} \neq 1$ in $(\mathbb{F}_q[t]/P^{p^{\ell+1}})^{(1)}$. As it is clear that $a^{p^\ell} \not\equiv 1 \pmod{P^{p^{\ell+1}}}$, the proof of the desired result is completed. \square

5.2. The Ext^1 -modules and t -modules. In this subsection, we will give an identification between certain Ext^1 -modules and t -modules. The key ingredient and ideas exhibited here are not new; actually they are due to G. Anderson, who shared his unpublished notes to the authors. Elements of these constructions are also present in the works of Hartl and Pink [HP04] and Taelman [Ta10]. In what follows, we fix two r -tuples $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$ and $\mathfrak{Q} \in \bar{k}[t]^r$ satisfying (2.3.1). Associated these two r -tuples, we let M (resp. M') be the Frobenius module defined by Φ as in (2.3.4) (resp. Φ' as in (2.3.5)).

Theorem 5.2.1. *Let $\{m_1, \dots, m_r\}$ be a $\bar{k}[t]$ -basis of M' on which the σ -action is presented by the matrix Φ' . Let $M \in \text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$ be defined by the matrix*

$$\begin{pmatrix} \Phi' & 0 \\ f_1, \dots, f_r & 1 \end{pmatrix}.$$

Then the map

$$\mu := (M \mapsto f_1 m_1 + \dots + f_r m_r) : \text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M') \rightarrow M' / (\sigma - 1)M'$$

is an isomorphism of $\mathbb{F}_q[t]$ -modules.

Proof. We first show that the map μ is well-defined. Suppose that M is trivial in $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M')$. Equivalently, there exists $u_1, \dots, u_r \in \bar{k}[t]$ so that

$$\begin{pmatrix} I_r & \\ u_1, \dots, u_r & 1 \end{pmatrix}^{(-1)} \begin{pmatrix} \Phi' & \\ f_1, \dots, f_r & 1 \end{pmatrix} = \begin{pmatrix} \Phi' & \\ & 1 \end{pmatrix} \begin{pmatrix} I_r & \\ u_1, \dots, u_r & 1 \end{pmatrix}.$$

The equation above is equivalent to

$$(f_1, \dots, f_r)(m_1, \dots, m_r)^{\text{tr}} = \left((u_1, \dots, u_r) - (u_1^{(-1)}, \dots, u_r^{(-1)})\Phi' \right) (m_1, \dots, m_r)^{\text{tr}},$$

which is equivalent to

$$(f_1, \dots, f_r)(m_1, \dots, m_r)^{\text{tr}} = (\sigma - 1) \left((-u_1, \dots, -u_r)(m_1, \dots, m_r)^{\text{tr}} \right).$$

So we have shown that μ is well-defined and also that μ is one to one. It is clear that μ is $\mathbb{F}_q[t]$ -linear and also surjective. \square

Now let us consider the n -th tensor power of the Carlitz module. The Frobenius module associated to $\mathbf{C}^{\otimes n}$ is the n -th tensor power of the Carlitz motive denoted by $\mathbf{C}^{\otimes n} := \bar{k}[t]$, on which σ acts by

$$\sigma(f) = (t - \theta)^n f^{(-1)}, \quad f \in \mathbf{C}^{\otimes n}.$$

As a $\bar{k}[\sigma]$ -module, $C^{\otimes n}$ is free of rank n with basis $\{(t - \theta)^{n-1}, \dots, t - \theta, 1\}$. From this observation, it is not hard to check that the Frobenius module M' fixed as above is a free left $\bar{k}[\sigma]$ -module of rank $d := (s_1 + \dots + s_r) + (s_2 + \dots + s_r) + \dots + s_r$, and

$$\left\{ (t - \theta)^{s_1 + \dots + s_r - 1} m_1, \dots, (t - \theta) m_1, m_1, \dots, (t - \theta)^{s_r - 1} m_r, \dots, (t - \theta) m_r, m_r \right\}$$

is a $\bar{k}[\sigma]$ -basis of M' . We further observe that $(t - \theta)^N M' / \sigma M' = (0)$ for $N \gg 0$ and hence M' is an Anderson t -motive in the sense of [Po8], which is called a dual t -motive in [ABPo4].

For such M' , we can identify $M' / (\sigma - 1)M'$ with the direct sum of d copies of \bar{k} as follows. Fixing a $\bar{k}[\sigma]$ -basis v_1, \dots, v_d of M' given as above, we can express any $m \in M'$ as

$$m = \sum_{i=1}^d u_i v_i, \quad u_i \in \bar{k}[\sigma],$$

and then we define $\Delta : M' \rightarrow \text{Mat}_{d \times 1}(\bar{k})$ by

$$(5.2.2) \quad \Delta(m) := \begin{pmatrix} \delta(u_1) \\ \vdots \\ \delta(u_d) \end{pmatrix},$$

where

$$\delta\left(\sum_i c_i \sigma^i = \sum_i \sigma^i c_i^{q^i}\right) = \sum_i c_i^{q^i}.$$

It follows that Δ is a morphism of \mathbb{F}_q -vector spaces with kernel $(\sigma - 1)M'$. We note that if $(a_1, \dots, a_d)^{\text{tr}} \in \text{Mat}_{d \times 1}(\bar{k})$, then there is a natural lift to M' , since

$$\Delta(a_1 v_1 + \dots + a_d v_d) = \begin{pmatrix} a_1 \\ \vdots \\ a_d \end{pmatrix}.$$

As $t(\sigma - 1)M' \subseteq (\sigma - 1)M'$, the map Δ induces an $\mathbb{F}_q[t]$ -module structure on $\text{Mat}_d(\bar{k})$. We denote by (E', ρ) the t -module defined over \bar{k} with $E'(\bar{k})$ identified with $\text{Mat}_d(\bar{k})$, on which the $\mathbb{F}_q[t]$ -module structure is given by

$$\rho : \mathbb{F}_q[t] \rightarrow \text{Mat}_d(\bar{k}[\tau])$$

so that

$$\Delta(t(a_1 v_1 + \dots + a_d v_d)) = \rho_t \begin{pmatrix} a_1 \\ \vdots \\ a_d \end{pmatrix}.$$

For example, we consider $C^{\otimes n}$. As a $\bar{k}[\sigma]$ -module, $C^{\otimes n}$ is free of rank n with basis $\{(t - \theta)^{n-1}, \dots, t - \theta, 1\}$. We let

$$\Delta_n : C^{\otimes n} \rightarrow \text{Mat}_{n \times 1}(\bar{k})$$

be defined as above with respect to this basis. For $(a_1, \dots, a_n)^{\text{tr}} \in \text{Mat}_{n \times 1}(\bar{k})$, we let

$$f = a_1(t - \theta)^{n-1} + \dots + a_{n-1}(t - \theta) + a_n,$$

so that $\Delta_n(f) = (a_1, \dots, a_n)^{\text{tr}}$. Now

$$tf = (\theta a_1 + a_2)(t - \theta)^{n-1} + \dots + (\theta a_{n-1} + a_n)(t - \theta) + (a_1^q + \theta a_n) + (\sigma - 1)(a_1^q),$$

and thus multiplication by t on $\text{Mat}_{n \times 1}(\bar{k})$ is given by

$$t \cdot \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix} = \Delta_n(tf) = [t]_n \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}.$$

In this way we identify $\mathbf{C}^{\otimes n} / (\sigma - 1)\mathbf{C}^{\otimes n}$ and $\mathbf{C}^{\otimes n}(\bar{k})$ as $\mathbb{F}_q[t]$ -modules. This identification between an abelian t -module over \bar{k} and the quotient of its associated dual t -motive modulo $\sigma - 1$ is due entirely to Anderson. One sees it in [ABP04, §4] (see especially the functor “ $f \pmod{\sigma - 1}$ ” in §4.1). See also [BP, §4.6], [CP11], [CP12] for other instances of this phenomenon for Drinfeld modules.

To summarize, we have the following result.

Theorem 5.2.3 (Anderson). *Let M' be the Frobenius module defined by the matrix Φ' in (2.3.5). Let (E', ρ) be the t -module with $E'(\bar{k})$ identified with $\text{Mat}_{d \times 1}(\bar{k})$, which is equipped with the $\mathbb{F}_q[t]$ -module structure via $\rho : \mathbb{F}_q[t] \rightarrow \text{Mat}_d(\bar{k}[\tau])$ through the map Δ as above. Then we have the following isomorphism of $\mathbb{F}_q[t]$ -modules*

$$M' / (\sigma - 1)M' \cong E'(\bar{k}),$$

and in fact E' is the t -module associated to the Anderson dual t -motive M' .

Remark 5.2.4. Combining the two theorems above we have that $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M') \cong E'(\bar{k})$ as $\mathbb{F}_q[t]$ -modules. Examples of this type of isomorphism were also studied by Ramachandran and the second author [PR03] for extensions of tensor powers of the Carlitz module. See also [S97, p. 529], [Ta10]. We further mention that in fact M' is a rigid analytically trivial Anderson t -motive as we have $\Psi'^{(-1)} = \Phi' \Psi'$ and so the corresponding t -module E' is uniformizable.

5.3. Reformulation of the criteria via t -modules.

Proposition 5.3.1. *Let n be a positive integer. Then for any nonzero polynomial $f \in A[t]$, we have $\Delta_n(f) \in \mathbf{C}^{\otimes n}(A)$. Equivalently, there exist $a_1, \dots, a_n \in A$ and $g \in \mathbf{C}^{\otimes n}$ so that*

$$f = a_1(t - \theta)^{n-1} + \dots + a_n + (\sigma - 1)g.$$

Proof. We write $f = \sum f_i \theta^i$ with $f_i \in \mathbb{F}_q[t]$. Then via the $\mathbb{F}_q[t]$ -linear map Δ_n we have

$$\Delta_n\left(\sum_i f_i \theta^i\right) = \sum_i [f_i]_n \Delta_n(\theta^i) = \sum_i [f_i]_n \begin{pmatrix} 0 \\ \vdots \\ \theta^i \end{pmatrix} \in \mathbf{C}^{\otimes n}(A).$$

□

Proposition 5.3.2. *Let M' be the Frobenius module defined by the matrix Φ' in (2.3.5) with a $\bar{k}[t]$ -basis m_1, \dots, m_r . Let $\{v_1, \dots, v_d\}$ be the $\bar{k}[\sigma]$ -basis of M' given by*

$$\left\{ (t - \theta)^{s_1 + \dots + s_r - 1} m_1, \dots, (t - \theta) m_1, m_1, \dots, (t - \theta)^{s_r - 1} m_r, \dots, (t - \theta) m_r, m_r \right\}.$$

Let Ξ be the set consisting of all elements in M' of the form $\sum_{i=1}^d e_i v_i$, where $e_j = \sum_n \sigma^n u_{nj}$ with each $u_{nj} \in A$. Then for any nonzero $f \in A[t]$ and any $1 \leq \ell \leq r$, we have that $f m_\ell \in \Xi$.

Proof. We first prove the case when $\ell = 1$. We divide f by $(t - \theta)^{s_1 + \dots + s_r}$ and write

$$f = g_1(t - \theta)^{s_1 + \dots + s_r} + \gamma_1,$$

where $g_1, \gamma_1 \in A[t]$ with $\deg_t \gamma_1 < s_1 + \dots + s_r$. So $fm_1 = g_1\sigma m_1 + \gamma_1 m_1 = \sigma g_1^{(1)} m_1 + \gamma_1 m_1$. Note that by expanding γ in terms of powers of $(t - \theta)$ we see that $\gamma_1 m_1$ is an A -linear combination of $\{\nu_1, \dots, \nu_{s_1 + \dots + s_r}\}$.

Next we divide $g_1^{(1)} \in A[t]$ by $(t - \theta)^{s_1 + \dots + s_r}$ and write

$$g_1^{(1)} = g_2(t - \theta)^{s_1 + \dots + s_r} + \gamma_2,$$

where $g_2, \gamma_2 \in A[t]$ with $\deg_t \gamma_2 < s_1 + \dots + s_r$. So

$$\sigma g_1^{(1)} m_1 = \sigma (g_2(t - \theta)^{s_1 + \dots + s_r} + \gamma_2) m_1 = \sigma^2 g_2^{(1)} m_1 + \sigma \gamma_2 m_2.$$

By expanding γ_2 in terms of powers of $(t - \theta)$ we see that $\sigma \gamma_2 m_2 \in \Xi$. By dividing $g_2^{(1)}$ by $(t - \theta)^{s_1 + \dots + s_r}$ and continuing the procedure as above inductively we eventually obtain that $fm_1 \in \Xi$.

Now for $\ell \geq 2$ we suppose that multiplication by any element of $A[t]$ on m_i belongs to Ξ for $1 \leq i \leq \ell - 1$. We prove that $fm_\ell \in \Xi$ by the induction on the degree of f in t , and note that the result is valid when $\deg_t f \leq s_\ell + \dots + s_r - 1$ by expanding f in terms of powers of $(t - \theta)$. So we suppose that $\deg_t f \geq s_\ell + \dots + s_r$.

We divide f by $(t - \theta)^{s_\ell + \dots + s_r}$ and write

$$f = g_1(t - \theta)^{s_\ell + \dots + s_r} + \gamma_1,$$

where $g_1, \gamma_1 \in A[t]$ with $\deg_t \gamma_1 < s_\ell + \dots + s_r$. It follows that

$$\begin{aligned} fm_\ell &= g_1(t - \theta)^{s_\ell + \dots + s_r} m_\ell + \gamma_1 m_\ell \\ &= g_1 \left\{ \sigma m_\ell - H_{s_{\ell-1}-1}^{(-1)}(t - \theta)^{s_{\ell-1} + \dots + s_r} m_{\ell-1} \right\} + \gamma_1 m_\ell \\ &= g_1 \left\{ \sigma m_\ell - \left\{ \sigma H_{s_{\ell-1}-1} m_{\ell-1} \right. \right. \\ &\quad \left. \left. - H_{s_{\ell-1}-1}^{(-1)} H_{s_{\ell-2}-1}^{(-1)}(t - \theta)^{s_{\ell-2} + \dots + s_r} m_{\ell-2} \right\} \right\} + \gamma_1 m_\ell \\ &\quad \vdots \\ &= g_1 \left\{ \sigma m_\ell + \sum_{i=1}^{\ell-1} (-1)^i \sigma \beta_1 \cdots \beta_i m_{\ell-i} \right\} + \gamma_1 m_\ell, \end{aligned}$$

where $\beta_i := H_{s_{\ell-i}-1} \in A[t]$ for $i = 1, \dots, \ell - 1$. It follows that

$$fm_\ell = \sigma g^{(1)} m_\ell + \sum_{i=1}^{\ell-1} (-1)^i \sigma g^{(1)} \beta_1 \cdots \beta_i m_{\ell-i} + \gamma_1 m_\ell.$$

However, by expanding γ_1 in terms of powers of $(t - \theta)$ we see that $\gamma_1 m_\ell \in \Xi$, and by hypothesis $\sum_{i=1}^{\ell-1} (-1)^i \sigma g^{(1)} \beta_1 \cdots \beta_i m_{\ell-i} \in \Xi$. Thus, to prove the desired result we are reduced to proving that $g^{(1)} m_\ell \in A[t]$, which is valid by the induction hypothesis since $\deg_t g^{(1)} = \deg_t g < \deg_t f$. \square

Remark 5.3.3. By (5.2.2) we see that $\Delta(\Xi) \subseteq E'(A)$.

Now we put $\Omega = (H_{s_1-1}, \dots, H_{s_r-1})$, where H_i are the Anderson-Thakur polynomials (see §4.1). We let $\mathbf{v}_s \in E'(\bar{k})$ be image of M under the composition of isomorphisms

$$\mathrm{Ext}_{\mathcal{F}}^1(\mathbf{1}, M') \cong M' / (\sigma - 1)M' \cong E'(\bar{k}).$$

Precisely,

$$\mathbf{v}_s := \Delta \left(H_{s_r-1}^{(-1)}(t - \theta)^{s_r} m_r \right).$$

Theorem 5.3.4. *For each r -tuple $\mathbf{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$, we have that*

- (a) *The associated t -module E' given above is defined over A ;*
- (b) *The point \mathbf{v}_s is an integral point in $E'(A)$.*

Proof. (a). Recall that M' is the Frobenius module defined by Φ' as in (2.3.5) with $\bar{k}[t]$ -basis m_1, \dots, m_r . Put $d = (s_1 + \dots + s_r) + \dots + s_r$ and let $\{v_1, \dots, v_d\}$ be the $\bar{k}[\sigma]$ -basis of M' given by

$$\left\{ (t - \theta)^{s_1 + \dots + s_r - 1} m_1, \dots, (t - \theta) m_1, m_1, \dots, (t - \theta)^{s_r - 1} m_r, \dots, (t - \theta) m_r, m_r \right\}.$$

We identify $M' / (\sigma - 1)M'$ with $\mathrm{Mat}_{d \times 1}(\bar{k})$ via the map Δ with respect to v_1, \dots, v_d .

Given any point $(a_1, \dots, a_d)^{\mathrm{tr}} \in E'(\bar{k})$, its corresponding element in $M' / (\sigma - 1)M'$ has a representative of the form $a_1 v_1 + \dots + a_d v_d$. We claim that the element

$$t \left(\sum_{i=1}^d a_i v_i \right)$$

can be expressed as $\sum_{i=1}^d b_i v_i \in \Xi$ for which each b_i is a of the form $b_i = \sum_j \sigma^j c_j$ so that c_j is an A -linear combination of $q^{(\cdot)}$ -th powers of the a'_h s. Then via the map Δ , the claim implies that the t -module E' is defined over A .

We observe that if some

$$v_i \notin \mathcal{S} := \left\{ (t - \theta)^{s_1 + \dots + s_r - 1} m_1, \dots, (t - \theta)^{s_{r-1} + s_r - 1} m_{r-1}, (t - \theta)^{s_r - 1} m_r \right\},$$

then

$$ta_i v_i = a_i(t - \theta)v_i + \theta a_i v_i = a_i v_{i-1} + \theta a_i v_i.$$

Therefore we reduce the claim to the case of $v_i \in \mathcal{S}$. To simplify the notation, we denote

$$v_{i_1} = (t - \theta)^{s_r - 1} m_r, \quad \dots, \quad v_{i_r} = (t - \theta)^{s_1 + \dots + s_r - 1} m_1.$$

Now given any $1 \leq \ell \leq r$ we consider $ta_{i_\ell} v_{i_\ell} = a_{i_\ell} t(t - \theta)^{s_1 + \dots + s_r - 1} m_\ell$. Applying Proposition 5.3.2 to $t(t - \theta)^{s_1 + \dots + s_r - 1} m_\ell$ we see that $ta_{i_\ell} v_{i_\ell}$ can be written as the form

$$a_{i_\ell} \sum_{j=1}^d \left(\sum_{e_j} \sigma^{e_j} b_{e_j} \right) v_j = \sum_{j=1}^d \left(\sum_{e_j} \sigma^{e_j} a_{i_\ell}^{q^{e_j}} b_{e_j} \right) v_j$$

for some $b_{e_j} \in A$, whence the desired result follows.

(b). Note that

$$\begin{aligned}
H_{s_r-1}^{(-1)}(t-\theta)^{s_r}m_r &= \sigma H_{s_r-1}m_r - H_{s_r-1}^{(-1)}H_{s_{r-1}-1}^{(-1)}(t-\theta)^{s_{r-1}+s_r}m_{r-1} \\
&\vdots \\
&= H_{s_r-1}^{(-1)}\left\{\sigma m_r + \sum_{i=1}^{r-1}(-1)^i\sigma\beta_1\cdots\beta_im_{r-i}\right\} \\
&= \sigma H_{s_r-1}m_r + \sum_{i=1}^{r-1}(-1)^i\sigma H_{s_r-1}\beta_1\cdots\beta_im_{r-i},
\end{aligned}$$

where $\beta_i := H_{s_{r-i}-1} \in A[t]$ for $i = 1, \dots, r-1$. Applying Proposition 5.3.2 to the right-hand side of the equation above we see that

$$H_{s_r-1}^{(-1)}(t-\theta)^{s_r}m_r \in \mathfrak{E}.$$

Since $\mathbf{v}_s = \Delta\left(H_{s_r-1}^{(-1)}(t-\theta)^{s_r}m_r\right)$, the result follows from Remark 5.3.3. \square

It follows that combining Theorems 4.2.2, 4.4.2, 5.2.1, and 5.2.3 we have the following criteria.

Theorem 5.3.5. *For any $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$, we have the following equivalence.*

- (1) $\zeta_A(s_1, \dots, s_r)$ is Eulerian.
- (2) \mathbf{v}_s is an $\mathbb{F}_q[t]$ -torsion point in the t -module $E'(A)$.

Finally Theorem 4.4.2 can now be transformed into the following concrete form:

Theorem 5.3.6. *Given $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$ with $w := \sum_{i=1}^r s_i$, put $\mathfrak{Q} := (H_{s_1-1}, \dots, H_{s_r-1})$ and $Q := H_{w-1}$, where $\{H_n\}$ be the Anderson-Thakur polynomials given in §4.1. Let Φ (resp. X) be defined as (2.3.4) (resp. (3.1.1)), and let M (resp. N) be the object in \mathcal{F} defined by Φ (resp. X). Let $M' \in \mathcal{F}$ be defined by Φ' given in (2.3.5) with a $\bar{k}[t]$ -basis $\{m_1, \dots, m_r\}$, and E' be the t -module associated to M' . Put $\mathbf{v}_s = \Delta\left(H_{s_r-1}^{(-1)}(t-\theta)^{s_r}m_r\right) \in E'(A)$ and $\mathbf{u}_s = \Delta\left(H_{w-1}^{(-1)}(t-\theta)^w m_1\right) \in E'(A)$. Then we have:*

- (a) *If w is not divisible by $q-1$, then we have that $\zeta_A(\mathfrak{s})$ is zeta-like if and only if there exists $a, b \in \mathbb{F}_q[t]$ (not both zero) so that $\rho_a(\mathbf{v}_s) + \rho_b(\mathbf{u}_s) = 0$ in the t -module $E'(A)$.*
- (b) *If w is divisible by $q-1$, then there exists nonzero $a \in \mathbb{F}_q[t]$ so that $\zeta_A(\mathfrak{s})$ is zeta-like if and only if $\rho_a(\mathbf{v}_s) = 0$ in the t -module $E'(A)$.*

Proof. Note that if w is not divisible by $q-1$, then $\tilde{\pi}^w \notin k_\infty$ and hence the k -linear dependence of $\{\zeta_A(\mathfrak{s}), \zeta_A(w), \tilde{\pi}^w\}$ is equivalent to that $\zeta_A(\mathfrak{s})/\zeta_A(w) \in k$. Thus, the result (a) follows from Theorem 4.4.2 and the identification $\text{Ext}_{\mathcal{F}}^1(\mathbf{1}, M') \cong E'(\bar{k})$. When w is divisible by $q-1$, we note that the zeta-like MZV's are the same as Eulerian MZV's because of (1.0.1), and hence the result (b) follows from Theorem 5.3.5. \square

6. THE ALGORITHM AND RULE SPECIFYING EULERIAN MZV'S

Fix the base finite field \mathbb{F}_q , as we are in positive characteristic p , multizeta values satisfy $\zeta_A(\mathfrak{s})^p = \zeta_A(s_1, \dots, s_r)^p = \zeta_A(ps_1, \dots, ps_r)$. Thus to investigate whether a given MZV is Eulerian we may restrict ourselves to consider only primitive tuples \mathfrak{s} , in the

sense that not all s_i are divisible by p . As first example of Eulerian MZV of depth > 1 , we cite e.g. Thakur [Togb, Thm. 5, Thm. 4]

$$\zeta_A(q-1, (q-1)^2) = \frac{1}{[1]_{q-1}} \zeta_A(q^2 - q),$$

where the Carlitz notation: $[\ell] := \theta^{q^\ell} - \theta$, is adopted, and the weight two Eulerian MZV

$$\zeta_A(q^\ell - 1, q^\ell(q-1)) = \zeta_A(q^\ell - 1) \zeta_A(q-1)^{q^\ell} - \zeta_A(q^{\ell+1} - 1).$$

This last relation has been extended inductively to arbitrary depth by Chen [Ch14], yielding Eulerian MZV of arbitrary depth r with respect to any \mathbb{F}_q . See (6.2.1).

Having Theorem 5.3.5 in our possession, we now write down an efficient algorithm for deciding whether any given MZV is Eulerian.

6.1. The algorithm. In accordance with Corollary 4.2.3, we only consider the case of all s_i divisible by $q-1$ when working on Eulerian MZVs. The following theorem offers an algorithm for Eulerian MZV's.

Theorem 6.1.1. *For any $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$ with all s_i divisible by $q-1$, we let*

$$w_i = s_{r-i} + s_{r-i+1} + \dots + s_r$$

for $i = 1, \dots, r-1$. We decompose

$$w_i = p^{\ell_i} n_i (q^{h_i} - 1)$$

so that $p \nmid n_i$ and h_i is the greatest integer for which $q^{h_i} - 1 \mid w_i$. Put

$$a = (t^{q^{h_{r-1}}} - t)^{p^{\ell_{r-1}}} \dots (t^{q^{h_1}} - t)^{p^{\ell_1}} \frac{\Gamma_{s_r+1}}{\Gamma_{s_r}} \text{den}(BC(s_r))|_{\theta=t}.$$

Then we have that $\zeta_A(s_1, \dots, s_r)$ is Eulerian if and only if $\rho_a(\mathbf{v}_\mathfrak{s}) = 0$.

Proof. Note that (\Leftarrow) follows from Theorem 5.3.5. We prove the result (\Rightarrow) by induction on the depth r . When $r = 1$, we write $\mathfrak{s} = s \in \mathbb{N}$. We claim that $\mathbf{v}_\mathfrak{s}$ is essentially the same as the special point Z_s in Definition 5.1.1, and so the result is valid by Remark 5.1.2. In this case, we note that $M' = \mathbb{C}^{\otimes s}$ and $\mathbf{v}_\mathfrak{s} = \Delta(H_{s-1}^{(-1)}(t-\theta)^s)$. We further note that

$$H_{s-1}^{(-1)}(t-\theta)^s \equiv H_{s-1} \pmod{(\sigma-1)},$$

which implies

$$\mathbf{v}_\mathfrak{s} := \Delta\left(H_{s-1}^{(-1)}(t-\theta)^s\right) = \Delta(H_{s-1}).$$

Let $H_{s-1} = \sum_{i \geq 0} h_{si} \theta^i$ with $h_{si} \in \mathbb{F}_q[t]$. Since the map Δ induces an $\mathbb{F}_q[t]$ -module isomorphism between $\mathbb{C}^{\otimes s} / (\sigma-1)\mathbb{C}^{\otimes s}$ and $\mathbb{C}^{\otimes s}(\bar{k})$, and Δ maps θ to the vector $(0, \dots, 0, \theta^i)^{\text{tr}} \in \mathbb{C}^{\otimes s}(A)$, we see that

$$\mathbf{v}_\mathfrak{s} = \Delta(H_{s-1}) = \Delta\left(\sum_{i \geq 0} h_{si} \theta^i\right) = \sum_{i \geq 0} [h_{si}]_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ \theta^i \end{pmatrix} = Z_s.$$

So the result is valid by Remark 5.1.2.

Suppose that the result is valid for depth less than r . Let Φ'' be the square matrix of size $r - 1$ cut from the right lower square of Φ' in (2.3.5), and let M'' be the Frobenius module defined by Φ'' . Therefore we have the exact sequence of Frobenius modules

$$0 \rightarrow \mathbf{C}^{\otimes(s_1+\cdots+s_r)} \rightarrow M' \rightarrow M'' \rightarrow 0.$$

Note that the snake lemma implies the exact sequence of $\mathbb{F}_q[t]$ -modules

$$0 \rightarrow \mathbf{C}^{\otimes(s_1+\cdots+s_r)} / (\sigma - 1)\mathbf{C}^{\otimes(s_1+\cdots+s_r)} \rightarrow M' / (\sigma - 1)M' \rightarrow M'' / (\sigma - 1)M'' \rightarrow 0$$

Denote by (E'', ϕ) the t -module underlying $M'' / (\sigma - 1)M''$, and so we have the exact sequence of $\mathbb{F}_q[t]$ -modules

$$0 \rightarrow \mathbf{C}^{\otimes(s_1+\cdots+s_r)}(\bar{k}) \rightarrow E'(\bar{k}) \rightarrow E''(\bar{k}) \rightarrow 0.$$

Denote by π the projection map $E'(\bar{k}) \rightarrow E''(\bar{k})$ given by

$$(a_1, \dots, a_n)^{\text{tr}} \mapsto (a_{s_1+\cdots+s_r+1}, \dots, a_r)^{\text{tr}}.$$

Put $\mathfrak{s} = (s_2, \dots, s_r)$. We claim that $\mathbf{v}_{\mathfrak{s}} = \pi(\mathbf{v}_{\mathfrak{s}})$. Assume this claim first. We write $a = (t^{q^{h_{r-1}}} - t)^{p^{\ell_{r-1}}} b$, where

$$b := (t^{q^{h_{r-2}}} - t)^{p^{\ell_{r-2}}} \cdots (t^{q^{h_1}} - t)^{p^{\ell_1}} \frac{\Gamma_{s_r+1}}{\Gamma_{s_r}} \text{den}(BC(s_r))|_{\theta=t},$$

then by the induction hypothesis we see that $\mathbf{0} = \phi_b(\mathbf{v}_{\mathfrak{s}}) = \pi(\rho_b(\mathbf{v}_{\mathfrak{s}}))$ and hence

$$\rho_b(\mathbf{v}_{\mathfrak{s}}) \in \text{Ker } \pi = \mathbf{C}^{\otimes(s_1+\cdots+s_r)}(\bar{k}).$$

Since by Theorem 5.3.4 E' and E'' are defined over A and $\mathbf{v}_{\mathfrak{s}}, \mathbf{v}_{\mathfrak{s}}$ are integral points, $\rho_b(\mathbf{v}_{\mathfrak{s}}) \in \mathbf{C}^{\otimes(s_1+\cdots+s_r)}(k)_{\text{tor}}$. Thus the result follows by Lemma 5.1.3.

Finally, we note that the claim above follows from the following commutative diagram

$$\begin{array}{ccc} M' & \xrightarrow{\Delta} & E'(\bar{k}) \\ \downarrow & & \downarrow \pi \\ M'' & \xrightarrow{\Delta} & E''(\bar{k}), \end{array}$$

where $M' \rightarrow M''$ is the projection map given by $\sum_{i=1}^r f_i m_i \mapsto \sum_{i=2}^r f_i m_i$ with $f_i \in \bar{k}[t]$. \square

For any $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^d$, let $\mathbf{u} = (u_1, \dots, u_r) \in (k^\times)^r \cap \mathbb{D}_{\mathfrak{s}}$ satisfy the hypotheses of Theorem 4.3.2. Applying the same arguments above we obtain the following result.

Corollary 6.1.2. *For any $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^d$ with all s_i divisible by $q - 1$, let $\mathbf{u} = (u_1, \dots, u_r) \in (k^\times)^r \cap \mathbb{D}_{\mathfrak{s}}$ satisfy the hypotheses of Theorem 4.3.2. Put $w_0 := s_r$ and write*

$$w_0 = p^{\ell_0} n_0 (q^{h_0} - 1)$$

so that $p \nmid n_0$ and h_0 is the greatest integer for which $q^{h_0} - 1 \mid w_0$. Let $(h_1, \ell_1), \dots, (h_{r-1}, \ell_{r-1})$ be defined in Theorem 6.1.1. Put $\mathfrak{Q} := \mathbf{u}$ and let M' be the Frobenius module defined by the matrix (2.3.5) with a $\bar{k}[t]$ -basis $\{m_1, \dots, m_r\}$. Let (E', ρ) be the t -module underlying associated to M' and $\mathbf{v}_{\mathfrak{s}} := \Delta \left(u_r^{(-1)} (t - \theta)^{s_r} m_r \right)$. Define $a := \prod_{i=0}^{r-1} (t^{q^{h_i}} - t)^{p^{\ell_i}} \in \mathbb{F}_q[t]$. Then we have that the value $\text{Li}_{\mathfrak{s}}(\mathbf{u})$ is Eulerian if and only if $\rho_a(\mathbf{v}_{\mathfrak{s}}) = 0$.

Proof. The proof is outlined as

- The t -module (E', ρ) is defined over k and \mathbf{v}_s is rational point in $E'(k)$ using the fact $\mathbf{u} \in (k^\times)^r \cap \mathbb{D}_s$ and following the arguments in Theorem 5.3.4.
- Note that for $r = 1$, we have $E' = \mathbf{C}^{\otimes s_r}$ from §5.2.
- In the case $r = 1$, we have that \mathbf{v}_s is an $\mathbb{F}_q[t]$ -torsion point in $E'(k)$ if and only if \mathbf{v}_s is $(t^{q^{h_0}} - t)^{p^{e_0}}$ -torsion by Lemma 5.1.3.
- The result follows by following the induction arguments in the proof of Theorem 6.1.1.

□

6.1.1. *The algorithm.* Here we provide the algorithm from Theorem 6.1.1. Given any $\mathfrak{s} = (s_1, \dots, s_r) \in \mathbb{N}^r$ with each s_i divisible by $q - 1$, we list the essential steps as follows.

- (I) Compute the Anderson-Thakur polynomials $H_{s_1-1}, \dots, H_{s_r-1}$.
- (II) Put $\tilde{\Omega} = (H_{s_1-1}, \dots, H_{s_r-1})$ and let M' be the Frobenius module defined by Φ' as in (2.3.5) with $\bar{k}[t]$ -basis m_1, \dots, m_r . Put $d = (s_1 + \dots + s_r) + \dots + s_r$ and let $\{v_1, \dots, v_d\}$ be the $\bar{k}[\sigma]$ -basis of M' given by

$$(t - \theta)^{s_1 + \dots + s_r - 1} m_1, \dots, (t - \theta) m_1, m_1, \dots, (t - \theta)^{s_r - 1} m_r, \dots, (t - \theta) m_r, m_r.$$

Identify $M' / (\sigma - 1)M'$ with $\text{Mat}_{d \times 1}(\bar{k})$ via v_1, \dots, v_d .

- (III) Write down the t -action on $M' / (\sigma - 1)M'$, and so giving a t -module structure on $\text{Mat}_{d \times 1}(\bar{k})$, which we denote by (E', ρ) .
- (IV) Consider $H_{s_r-1}^{(-1)}(t - \theta)^{s_r} m_r \in M' / (\sigma - 1)M'$, which corresponds to an integral point $\mathbf{v}_s = (a_1, \dots, a_d)^{\text{tr}} \in E'(A)$ from the decomposition $H_{s_r-1}^{(-1)}(t - \theta)^{s_r} m_r \equiv \sum_{i=1}^d a_i v_i \pmod{\sigma - 1}$.
- (V) Define the polynomial a as in Theorem 6.1.1, and then compute $\rho_a(\mathbf{v}_s)$. If it is zero, then $\zeta_A(s_1, \dots, s_r)$ is Eulerian; otherwise, $\zeta_A(s_1, \dots, s_r)$ is non-Eulerian.

6.1.2. *Examples of (E', ρ) and \mathbf{v}_s .* We provide some examples of the explicit forms of (E', \mathbf{v}_s) . The following are two examples associated to Eulerian MZV's, i.e., \mathbf{v}_s is $\mathbb{F}_q[t]$ -torsion in $E'(A)$.

- (1) Let $q = 3, \mathfrak{s} = (2, 4)$. Then (E', ρ) associated to $\zeta(2, 4)$ is given by

$$\rho_t = \begin{pmatrix} \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 \\ \tau & 0 & 0 & 0 & 0 & \theta & 2\tau & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & \tau & 0 & 0 & \theta \end{pmatrix}$$

and

$$\mathbf{v}_s = (0, 0, 1, 0, 1, (\theta + 2\theta^3), 2, 0, 2, (2\theta + \theta^3))^{\text{tr}}.$$

(2) Let $q = 2$, $\mathfrak{s} = (1, 2, 4)$. Then (E', ρ) associated to $\zeta(1, 2, 4)$ is given by

$$\rho_t = \begin{pmatrix} \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \tau & 0 & 0 & 0 & 0 & 0 & \theta & \tau & 0 & 0 & 0 & 0 & 0 & \tau & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \tau & 0 & \theta \end{pmatrix}$$

and

$$\mathbf{v}_{\mathfrak{s}} = (0, 0, 0, 0, 1, 1, (\theta + \theta^2), 0, 0, 0, 1, 1, (\theta + \theta^2), 0, 1, 1, (\theta + \theta^2))^{\text{tr}}.$$

The following are two examples associated to non-Eulerian MZV's, $\mathbf{v}_{\mathfrak{s}}$ is not $\mathbb{F}_q[t]$ -torsion in $E'(A)$.

(3) Let $q = 3$, $\mathfrak{s} = (4, 2)$. Then (E', ρ) associated to $\zeta(4, 2)$ is given by

$$\rho_t = \begin{pmatrix} \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \theta & 1 & 0 & 0 & \tau & 0 & 0 \\ 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \theta & 1 & \tau & 0 & 0 \\ \tau & 0 & 0 & 0 & 0 & \theta & (\theta + 2\theta^3)\tau & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \tau & 0 & \theta \end{pmatrix}$$

and

$$\mathbf{v}_{\mathfrak{s}} = (0, 0, 1, 0, 1, (\theta + 2\theta^3), 0, 1)^{\text{tr}}.$$

(4) Let $q = 3$, $\mathfrak{s} = (2, 2, 2)$. Then (E', ρ) associated to $\zeta(2, 2, 2)$ is given by

$$\rho_t = \begin{pmatrix} \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 & 0 \\ \tau & 0 & 0 & 0 & 0 & \theta & 2\tau & 0 & 0 & 0 & \tau & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \tau & 0 & 0 & \theta & 2\tau & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \theta & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \tau & \theta \end{pmatrix}$$

and

$$\mathbf{v}_{\mathfrak{s}} = (0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 2, 0, 1)^{\text{tr}}.$$

6.2. Searching for the rules governing Eulerian MZV's. Lara Rodríguez and Thakur [LRT14] have given conjectures on which r -tuples (s_1, \dots, s_r) may occur for Eulerian MZV's. Furthermore, they also provided conjectural formulas for these special values. Computations based on implementing the above algorithm in Magma (by Yi-Hsuan Lin) have led us to the following description of Eulerian MZV's in arbitrary depth.

Fix prime power q , and call the sequence of r -tuples below Eulerian r -tuples with respect to \mathbb{F}_q :

$$\text{Euler}_1 := (q - 1) \text{ and } \text{Euler}_{r+1} := (q - 1, q \text{Euler}_r) \in \mathbb{N}^{r+1}.$$

For each depth r , we introduce a sequence of r -tuples in \mathbb{N}^r as follows:

$$\text{Euler}_r(\ell) := (q^\ell - 1, q^\ell \text{Euler}_{r-1}), \text{ for } r > 1, \ell \geq 1,$$

and $\text{Euler}_1(\ell) := (q^\ell - 1)$. Call this the canonical sequence of depth r with respect to \mathbb{F}_q . The corresponding MZV's $\zeta_A(\text{Euler}_r(\ell))$ are all Eulerian. This follows from the Euler-Carlitz formula(1.0.1) and the following inductive formula of Chen [Ch14] for all $r \geq 2$ and $\ell \geq 1$:

$$(6.2.1) \quad \zeta_A(\text{Euler}_r(\ell)) = \zeta_A(q^\ell - 1)\zeta_A(\text{Euler}_{r-1})^{q^\ell} - \zeta_A(\text{Euler}_{r-1}(\ell + 1)).$$

Note that when $q = 2$, all depth one Carlitz zeta values $\zeta_A(n)$ are Eulerian, and $\zeta_A(\mathfrak{s})$ is Eulerian if and only if it is zeta-like. The following Eulerian multizeta values of arbitrary depth $r > 1$ and weight 2^{r-1} have been found by Lara Rodríguez and Thakur [LRT14]:

$$(6.2.2) \quad \zeta_A(1, \mathfrak{s}) := \zeta_A(1, 1, 2, \dots, 2^{r-1}) = \frac{1}{[1]^{2^{r-1}}[2]^{2^{r-2}} \dots [r]} \zeta_A(2^{r-1}).$$

For $q \geq 3$, we predict that the primitive Eulerian MZV's of depth $r \geq 2$ are precisely:

- (1) The canonical family with every depth $r \geq 2$, $\ell \geq 1$,

$$\zeta_A(\text{Euler}_r(\ell)), \text{ of weight } q^{r+\ell-1} - 1,$$

- (2) An extra family in depth $r = 2$, $\ell \geq 1$,

$$\zeta_A(q^\ell(q-1), q^{\ell+2} - 1 - q^\ell(q-1)), \text{ of weight } q^{\ell+2} - 1.$$

- (3) An exceptional primitive Eulerian MZV in depth $r = 2$,

$$\zeta_A(q-1, (q-1)^2), \text{ of weight } q^2 - q.$$

Thus for $q \geq 3$ there should exist depth $r > 1$ primitive Eulerian MZV's only in weights $q^2 - q$ (depth 2), and $q^\ell - 1$ (in any depth) for $\ell \geq r$. For depth $r = 2$, only in weights $q^2 - q$, $q^2 - 1$, and each weight has only one primitive Eulerian MZV. For weight $q^\ell - 1$, $\ell \geq 3$, each weight has two primitive Eulerian MZV's, coming from the two families in (1) and (2). Lara Rodríguez and Thakur [LRT14] have also given precise formulas (valid for any q) for the family (2):

$$\zeta_A(q^\ell(q-1), q^{\ell+2} - 1 - q^\ell(q-1)) = \frac{1}{[1]^{q^\ell(q-1)}} \zeta_A(q^{\ell+2} - 1).$$

In the case $q = 2$, we predict that the primitive Eulerian MZV's are given by:

- (1) The canonical family with every depth $r \geq 2$, $\ell \geq 1$,

$$\zeta_A(\text{Euler}_r(\ell)), \text{ of weight } 2^{r+\ell-1} - 1.$$

- (2) The extra family in depth $r = 2$, $\ell \geq 1$,

$$\zeta_A(2^\ell, 2^{\ell+2} - 1 - 2^\ell) \text{ weight } 2^{\ell+2} - 1.$$

- (3) Three exceptional primitive Eulerian MZV in depth $r = 2$:

$$\zeta_A(1, 1), \zeta_A(1, 3) = \zeta_A(1, 2^r - 1) = \left(\frac{1}{[1][2]} + \frac{1}{[1]} \right) \zeta_A(4), \text{ and}$$

$$\zeta_A(3, 5) = \frac{[2]^2 + 1}{[1]^4[2]} \zeta_A(8).$$

Thus primitive Eulerian pairs exist only in weights, 2, 3, 4, 7, 8, $2^\ell - 1$, $\ell \geq 4$. For each weight $2^\ell - 1$, $\ell \geq 3$, there are exactly two primitive Eulerian pairs from the two families. For weights 2, 3, 4, 8, each weight has only one primitive Eulerian pair.

- (4) There are exceptional primitive Eulerian MZV's for depth $r > 2$: $\zeta_A(1, \mathfrak{s})$, where \mathfrak{s} is a primitive Eulerian tuple of depth $r - 1$ and weight either 2^r or 2^{r-1} . Thus for depth $r > 3$, in each weight $2^r, 2^{r-1}$ there is only one exceptional primitive Eulerian MZV. In the case of depth 3, there are two exceptional primitive Eulerian of weight 8, and one exceptional primitive Eulerian of weight 4.

The exceptional sequence of primitive Eulerian MZV's of weight 2^{r-1} , $r > 1$, is the one given in (6.2.2). That the above exceptional sequence of primitive MZV's of weight 2^r consists only of Eulerians (first conjectured by Lara Rodríguez and Thakur [LRT14]) is a consequence of the following formula of Chen [Ch14]:

$$\zeta_A(1, \mathfrak{s}) = \zeta_A(1, 3, 2^2, \dots, 2^{r-1}) = \zeta_A(1)\zeta_A(1, 2, \dots, 2^{r-1}) + \zeta_A(1, 1, 2, \dots, 2^{r-2})^2.$$

When $r = 2$, this last equality also goes back to Thakur [To9b, Thm. 8]

All MZV's in the above list have been confirmed to be Eulerian by [Ch14] and [LRT14]. Our computations suggest that the above list exhausts all primitive Eulerian multizeta values for $\mathbb{F}_q[\theta]$. In other words, any r -tuple \mathfrak{s} of depth $r > 1$, not accounted by our list above should give non-Eulerian $\zeta_A(\mathfrak{s})$. Previously in [LRT14], Lara Rodríguez and Thakur had also collected data basing on continued fraction computations to decipher the occurrence of Eulerian MZV's, and made precise conjectures characterizing Eulerian tuples. Their conjectures agree with the above list. Our “ t -motivic” algorithm for determining Eulerian multizeta values is rooted by an entirely different principle, runs a bit more efficiently and is completely algebraic. It allows us to do computations inductively for higher depth because of the key Corollary 4.2.3, thereby leading to the above description which we believe is a complete list.

Summary of data certified by our computations

All tuples \mathfrak{s} of depth r and weights w within the following respective bounds have been checked for the Eulerian property. The answers agree with the description above and the MZV's in the complementary part of the list above are non-Eulerian. When $3 \leq q \leq 11$, all tuples having their weights within the bounds below have been checked, with no restriction on their depths except $q = 2$.

$$q = 2, \text{ depth} = 2, \text{ weight} \leq 256$$

$$q = 2, \text{ depth} = 3, 4, 5, \text{ weight} \leq 128$$

$$q = 2, \text{ depth} = 6, \text{ weight} \leq 64$$

$$q = 3, \text{ weight} \leq 243$$

$$q = 4, \text{ weight} \leq 256$$

$$q = 5, 7 \text{ weight} \leq q^3$$

$$8 \leq q \leq 19, \text{ weight} \leq q^2$$

REFERENCES

- [A86] G. W. Anderson, *t-motives*, Duke Math. J. **53** (1986), no. 2, 457–502.
 [ABPo4] G. W. Anderson, W. D. Brownawell and M. A. Papanikolas, *Determination of the algebraic relations among special Γ -values in positive characteristic*, Ann. of Math. (2) **160** (2004), no. 1, 237–313.
 [AT90] G. W. Anderson and D. S. Thakur, *Tensor powers of the Carlitz module and zeta values*, Ann. of Math. (2) **132** (1990), no. 1, 159–191.
 [AT09] G. W. Anderson and D. S. Thakur, *Multizeta values for $\mathbb{F}_q[t]$, their period interpretation, and relations between them*, Int. Math. Res. Not. IMRN (2009), no. 11, 2038–2055.

- [B12a] F. Brown, *On the decomposition of motivic multiple zeta values*, in: Galois-Teichmüller Theory and Arithmetic Geometry, Adv. Stud. Pure Math. **63**, Math. Soc. Japan, Tokyo, 2012, pp. 31–58.
- [B12b] F. Brown, *Mixed Tate motives over \mathbb{Z}* , Ann. of Math. (2) **175** (2012), no. 2, 949–976.
- [BP] W. D. Brownawell and M. A. Papanikolas, *A rapid introduction to Drinfeld modules, t -modules, and t -motives*, available at <http://www.math.tamu.edu/~map/BanffSurvey.pdf>, 2011.
- [Ca35] L. Carlitz, *On certain functions connected with polynomials in a Galois field*, Duke Math. J. **1** (1935), no. 2, 137–168.
- [C14] C.-Y. Chang, *Linear independence of monomials of multizeta values in positive characteristic*, to appear in Compos. Math., DOI: <http://dx.doi.org/10.1112/S0010437X1400743X>.
- [CP11] C.-Y. Chang and M. A. Papanikolas, *Algebraic relations among periods and logarithms of rank 2 Drinfeld modules*, Amer. J. Math. **133** (2011), no. 2, 359–391.
- [CP12] C.-Y. Chang and M. A. Papanikolas, *Algebraic independence of periods and logarithms of Drinfeld modules. With an appendix by Brian Conrad*, J. Amer. Math. Soc. **25** (2012), no. 1, 123–150.
- [CY07] C.-Y. Chang and J. Yu, *Determination of algebraic relations among special zeta values in positive characteristic*, Adv. Math. **216** (2007), no. 1, 321–345.
- [Ch14] H.-J. Chen, *On Eulerian multizeta values for $\mathbb{F}_q[t]$* , in preparation.
- [HP04] U. Hartl and R. Pink, *Vector bundles with a Frobenius structure on the punctured unit disc*, Compos. Math. **140** (2004), no. 3, 689–716.
- [G096] D. Goss, *Basic structures of function field arithmetic*, Springer-Verlag, Berlin, 1996.
- [LRT14] J. A. Lara Rodríguez and D. S. Thakur, *Zeta-like multizeta values for $\mathbb{F}_q[t]$* , Indian J. Pure Appl. Math. **45** (5), 785–798 (2014).
- [Po8] M. A. Papanikolas, *Tannakian duality for Anderson-Drinfeld motives and algebraic independence of Carlitz logarithms*, Invent. Math. **171** (2008), no. 1, 123–174.
- [PR03] M. A. Papanikolas and N. Ramachandran, *A Weil-Barsotti formula for Drinfeld modules*, J. Number Theory **98** (2003), no. 2, 407–431.
- [S97] S. K. Sinha, *Periods of t -motives and transcendence*, Duke Math. J. **88** (1997), no. 3, 465–535.
- [Ta10] L. Taelman, *1- t -motifs*, arXiv:0908.1503, 2010.
- [To4] D. S. Thakur, *Function field arithmetic*, World Scientific Publishing, River Edge NJ, 2004.
- [To9a] D. S. Thakur, *Power sums with applications to multizeta and zeta zero distribution for $\mathbb{F}_q[t]$* , Finite Fields Appl. **15** (2009), no. 4, 534–552.
- [To9b] D. S. Thakur, *Relations between multizeta values for $\mathbb{F}_q[T]$* , Int. Math. Res. Notices IMRN (2009), no. 12, 2318–2346.
- [Yu91] J. Yu, *Transcendence and special zeta values in characteristic p* , Ann. of Math. (2) **134** (1991), no. 1, 1–23.
- [Yu97] J. Yu, *Analytic homomorphisms into Drinfeld modules*, Ann. of Math. (2) **145** (1997), no. 2, 215–233.

DEPARTMENT OF MATHEMATICS, NATIONAL TSING HUA UNIVERSITY AND NATIONAL CENTER FOR THEORETICAL SCIENCES, HSINCHU CITY 30042, TAIWAN R.O.C.

E-mail address: cychang@math.nthu.edu.tw

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, U.S.A.

E-mail address: map@math.tamu.edu

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY AND TIMS, TAIPEI CITY 106, TAIWAN R.O.C.

E-mail address: yu@math.ntu.edu.tw